

Analysis of a Ferric Uptake Regulator (Fur) Mutant of *Desulfovibrio vulgaris* Hildenborough[∇]

Kelly S. Bender,^{1,10}† Huei-Che Bill Yen,^{1,10} Christopher L. Hemme,^{2,3,10} Zamin Yang,^{2,10}
Zhili He,^{2,3,10} Qiang He,^{4,10} Jizhong Zhou,^{2,3,10} Katherine H. Huang,^{5,10} Eric J. Alm,^{5,6,10}
Terry C. Hazen,^{7,10} Adam P. Arkin,^{5,8,9,10} and Judy D. Wall^{1,10*}

Department of Biochemistry, University of Missouri—Columbia, Columbia, Missouri 65211¹; Environmental Sciences Division, Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831²; Institute for Environmental Genomics, Department of Botany and Microbiology, University of Oklahoma, Norman, Oklahoma 73019³; Department of Civil and Environmental Engineering, Temple University, Philadelphia, Pennsylvania 19122⁴; Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720⁵; Biological Engineering Division, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139⁶; Earth Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720⁷; Department of Bioengineering, University of California, Berkeley, California 94720⁸; Howard Hughes Medical Institute, Chevy Chase, Maryland 20815⁹; and Virtual Institute for Microbial Stress and Survival¹⁰‡

Received 2 February 2007/Accepted 12 June 2007

Previous experiments examining the transcriptional profile of the anaerobe *Desulfovibrio vulgaris* demonstrated up-regulation of the Fur regulon in response to various environmental stressors. To test the involvement of Fur in the growth response and transcriptional regulation of *D. vulgaris*, a targeted mutagenesis procedure was used for deleting the *fur* gene. Growth of the resulting Δfur mutant (JW707) was not affected by iron availability, but the mutant did exhibit increased sensitivity to nitrite and osmotic stresses compared to the wild type. Transcriptional profiling of JW707 indicated that iron-bound Fur acts as a traditional repressor for ferrous iron uptake genes (*feoAB*) and other genes containing a predicted Fur binding site within their promoter. Despite the apparent lack of siderophore biosynthesis genes within the *D. vulgaris* genome, a large 12-gene operon encoding orthologs to TonB and TolQR also appeared to be repressed by iron-bound Fur. While other genes predicted to be involved in iron homeostasis were unaffected by the presence or absence of Fur, alternative expression patterns that could be interpreted as repression or activation by iron-free Fur were observed. Both the physiological and transcriptional data implicate a global regulatory role for Fur in the sulfate-reducing bacterium *D. vulgaris*.

Iron is an essential nutrient for most bacteria because of its role as an enzymatic cofactor and electron transport protein component. In addition to the metabolic importance of iron, pathogenic bacteria use its availability as an environmental signal for regulation of virulence genes. Despite the metabolic dependence on iron, cellular concentrations must be intricately regulated in aerobic environments to prevent Fe(II)-mediated formation of reactive oxygen species via Fenton chemistry (72). In most bacteria, this complex regulation is carried out by the ferric uptake regulator protein (Fur) (23, 29). The traditional mode of Fur regulation has been described as follows: under iron-replete conditions, the Fur protein and its corepressor [Fe(II)] block the transcription of iron uptake and storage genes. When iron becomes limiting, the Fur repressor is no longer saturated with Fe(II) and cannot bind the operator, leading to transcription of genes involved in iron uptake and storage. Since iron is found predominantly as insoluble ferric hydroxides in aerobic environments, bacteria have evolved a

mechanism for uptake dependent on the synthesis and transport of specialized chelators called siderophores. Thus, Fur also regulates the synthesis of siderophores in bacteria studied to date (29, 65, 68). In addition to its role as the primary regulator responding to available iron, Fur has also been shown to play a global regulatory role in oxidative stress response, acid tolerance response, virulence factor synthesis, and motility (2, 23, 29).

Because of the insolubility and sequestration of iron in aerobic and host environments, studies involving iron regulation have focused primarily on aerobic/facultative and pathogenic microbes, respectively. The role of Fur or iron metabolism in strict anaerobes has received little attention. In anaerobic environments, enough iron is expected to be in the Fe(II) form that its accessibility should not be limiting. Also, the likelihood of Fe(II)-mediated formation of reactive oxygen species via Fenton chemistry is decreased by limited oxygen exposure. Thus, elaborate iron regulation would seem to be less critical in anaerobes. However, recent genomic studies of metal-reducing δ -proteobacteria have indicated the presence of not one but three *fur* paralogs, *fur* (DVU0942), *perR* (DVU3095), and *zur* (DVU1340), in these anaerobes (61). While the roles of these regulators remain unclear, the sulfate reducers *Desulfovibrio vulgaris* Hildenborough and *Desulfovibrio desulfuricans* G20 appear to have extended Fur regulons compared to the predictions from genomes of other δ -proteobacteria (61). Regulon

* Corresponding author. Mailing address: Department of Biochemistry, 117 Schweitzer Hall, Columbia, MO 65211. Phone: (573) 882-8726. Fax: (573) 882-5635. E-mail: wallj@missouri.edu.

† Present address: Department of Microbiology, Southern Illinois University, Carbondale, IL 62901.

‡ <http://vimss.lbl.gov>.

[∇] Published ahead of print on 13 July 2007.

members include ferrous iron transporter genes (*feoAB*, DVU2572/71), a flavodoxin gene (*fld*, DVU2680), genes for P-type and ABC ATPases, and genes possessing GGDEF and HD domains. The proteins encoded by the latter genes are predicted to have cyclic di-GMP synthesis and hydrolysis activity, respectively, that could allow second-messenger concentrations to be responsive to Fur signals (66). A large cluster of genes predicted to be involved in biopolymer transport, such as *tonB* (DVU2390), were also suggested to be part of the Fur regulon (61). However, siderophore production by *Desulfovibrio* has not been documented, nor have genes for siderophore synthesis been identified. The annotation of a putative transporter for the siderophore enterobactin (*fepC*, DVU0648) suggests a possible mechanism for acquiring insoluble iron via chelators produced by other bacteria.

Desulfovibrio species are anaerobic sulfate-reducing bacteria (SRB) known for their ability to corrode ferrous metals as well as to reduce heavy metals such as uranium(VI), chromium(VI), and technetium(VII). *D. vulgaris* Hildenborough is also believed to possess a robust iron requirement based on its abundance of iron-containing cytochromes, hydrogenases, and electron transport proteins, as determined from genome analysis (36). Despite the solubility of iron in anaerobic environments, the degree to which Fe(II) is accessible to sulfate reducers is unknown, especially since sulfide, a by-product of sulfate reduction, complexes with Fe(II) to form insoluble pyrite (FeS). Another notable aspect of iron metabolism in *Desulfovibrio* is the production of a ferritin protein containing a unique heme group (58, 62). Ferritin and bacterioferritin proteins are produced by aerobic bacteria to sequester free iron in a nonreactive insoluble Fe(III) form (1, 2, 72). The role of these proteins in anaerobic bacteria has been assumed to be in oxygen defense (60, 62), but their intrinsic regulatory mechanisms are currently unknown.

Even though *Desulfovibrio* species are considered strict anaerobes, they do possess numerous oxygen detoxification proteins, such as rubrerythrin, rubredoxin oxidoreductase, and superoxide dismutase, all of which contain iron (19, 24, 45). The role of iron in response to oxygen stress was underscored by recent research with *Bacillus subtilis* indicating that iron oxidation was likely the initiating event for detection of peroxides via the metal-dependent peroxide sensor/regulator PerR (41). The annotation of a *perR* gene in SRB suggests that the relationship between these anaerobes and iron may be more complex than previously considered. Here we describe a marker exchange method for gene deletion in *D. vulgaris* Hildenborough and the physiology and transcriptional profile of the resulting Δfur strain. The transcriptional data are strengthened by comparison with previous studies documenting the differentially expressed genes in stressed *D. vulgaris* cells (12, 35, 46, 47). While Fur in *D. vulgaris* appears to regulate genes traditionally involved in Fe(II) uptake, our data suggest a diverse regulatory pattern for Fur in SRB. To our knowledge, this is the first report of a Δfur deletion in a sulfate-reducing bacterium.

MATERIALS AND METHODS

Bacterial strains, plasmids, and media. Bacterial strains and vectors used in this study are listed in Table 1. *D. vulgaris* cultures were grown anaerobically as previously described (57) in two defined media: LS4D, routinely used in transcriptomic and proteomic studies of this bacterium so data can be compared, and

TABLE 1. Bacteria and plasmids used in this study

Strain or plasmid	Description of relevant features	Reference or source
Strains		
<i>D. vulgaris</i> Hildenborough	Wild type; NCIMB 8303	36, 53
<i>D. vulgaris</i> JW707	Δfur Km ^r	This study
<i>E. coli</i> JM109	Cloning strain	Promega
Plasmids		
pBluescript SK(+)	Blue-white cloning vector; Amp ^r	Stratagene
pSC27	<i>Desulfovibrio</i> shuttle vector; source of Km ^r cassette	63
pMO707	pBluescript containing 2,839-kb <i>fur</i> deletion cassette; Amp ^r , Km ^r	This study

Yen45, used to reduce the formation of precipitates when ion concentrations were altered. LS4D (60 μ M total iron) consisted of 60 mM sodium lactate, 50 mM Na₂SO₄, 8.0 mM MgCl₂, 20 mM NH₄Cl, 2.2 mM K₂HPO₄, 0.6 mM CaCl₂, 30 mM PIPES [piperazine-N,N'-bis(2-ethanesulfonic acid)], 12.5 ml trace mineral solution per liter (7), NaOH to a pH of 7.2, and 1.0 ml 10 \times vitamin solution per liter (7) (added after autoclaving). Yen45 (30 μ M total iron) consisted of 60 mM sodium lactate, 30 mM Na₂SO₄, 8.0 mM MgCl₂, 20 mM NH₄Cl, 2 mM phosphate buffer (K₂HPO₄/NaH₂PO₄, pH 7.4), 0.6 mM CaCl₂, 30 mM Tris-HCl (pH 7.4), and, per liter, 2 ml modified trace mineral solution (7) (modified by omission of nitrilotriacetic acid and FeCl₂), 0.24 ml iron solution (125 mM FeCl₂/250 mM EDTA, pH 7.3), 20 ml sterile 1 M NaHCO₃ (added after autoclaving), and 1.0 ml sterile 10 \times vitamin solution (7) (added after autoclaving). As the reductant for both media, 5 ml per liter of an anaerobic titanium citrate solution was used. This solution contained 20% (wt/vol) titanium(III) chloride, 0.2 M sodium citrate, and 8.0% (wt/vol) sodium carbonate. For plating, LS4D medium was supplemented with 0.2% (wt/vol) yeast extract, 1.5% (wt/vol) agar, and 1.2 mM thioglycolate. Cells were distributed in 4 ml of molten top agar (30 mM PIPES, 1.5% agar).

Deletion cassette construction. Deletion cassettes were constructed by a method similar to the molecular bar-coding methods described for *Saccharomyces cerevisiae* (27, 69). Briefly, PCR primer sets were designed to amplify approximately 800 bp up- and downstream of the *fur* open reading frame (ORF) (DVU0942). Primer nucleotide sequences are listed in Table 2 along with genome locations relative to the *fur* ORF. For future tracking of the mutant in a mixed population, unique barcode sequences were added between the common sequences and Km^r sequences of primers P5 and P6. The following PCR mixture was used to amplify the upstream (primers P1 and P2) and downstream (primers P3 and P4) regions flanking the *fur* ORF from *D. vulgaris* as well as the Km^r cassette from pSC27 (63) (primers P5 and P6): 1 \times Hercules buffer (Stratagene, La Jolla, CA), 0.25 mM each deoxynucleoside triphosphate, 2.5 pmol each primer, 2.5 units Hercules polymerase (Stratagene), and 1 μ l template DNA in a total volume of 40 μ l. Reaction mixtures were cycled according to the following program: 94°C denaturation for 60 s followed by 5 cycles consisting of 94°C for 30 s, 45°C for 30 s, and 70°C for 70 s and then 24 cycles consisting of 94°C for 30 s, 54°C for 30 s, and 70°C for 70 s, ending with a 6-min extension at 70°C. Products were gel purified using Quantum Prep Freeze'N Squeeze DNA gel extraction spin columns (Bio-Rad, Hercules, CA) and immediately used for fusion PCR.

For the fusion PCR, the following reaction mixture was used: 1 \times Hercules buffer, 0.25 mM each deoxynucleoside triphosphate, 4 units Hercules polymerase, and 50 to 100 ng of each of the three products generated in the PCRs described above (approximately 3 μ l each). Before addition of the polymerase, the reaction mixture was heated to 94°C for 30 s. Reaction mixtures were subjected to four cycles consisting of 94°C for 30 s, 55°C for 60 s, 72°C for 3 min 30 s, and then 4 U more of polymerase and 25 pmol each of primers P1 and P4 were added. This mixture was then cycled 25 times at 94°C for 30 s, 55°C for 60 s, and 72°C for 3 min 30 s, ending with an 8-min extension at 72°C for 8 min. The resulting fusion PCR product was gel purified as described above and ligated into the EcoRV site of pBluescript SK(+) (Stratagene, La Jolla, CA) to generate pMO707. Prior to electroporation of pMO707 into *D. vulgaris*, the construct was sequenced to check for proper arrangement and lack of sequence errors.

Transformation and mutant selection. Approximately 10⁹ to 10¹⁰ *D. vulgaris* cells, harvested at early stationary phase (optical density at 600 nm [OD₆₀₀] of ca.

TABLE 2. PCR primers for deletion cassette construction

Primer	Sequence (5'-3') ^a	Position of 5' end
P1	<u>CTCTCTGCAACCTGACGGCG</u>	-780 bp relative to <i>fur</i> ORF
P2	<u>AAGACTGTAGCCGTACCTCGAATCTACTCTTGGGTTGGTGTCTGTGC</u>	-13 bp relative to <i>fur</i> ORF
P3	<u>AATCCGCTCACTAAGTTCATAGACCGAAGCGCATGTGAAGACTGCTGT</u>	+5 bp relative to <i>fur</i> ORF
P4	<u>GCACGAAGGACTCGAGAGGATTTCGATGC</u>	+835 bp relative to <i>fur</i> ORF
P5	<u>TAGATTCGAGGTACGGCTACAGTCTTGTGGACTGACGGCTAATCTCCCCAGAGT</u> CCCGCTCAG	+14 bp relative to Km ^r ORF
P6	<u>CGGTCTATGAACTTAGTGAGCGGATTTCGTAACGGCTTGTACCCAGAGAGGTAGCT</u> TGCAGTGGGCT	-205 bp relative to Km ^r ORF
P7	<u>GTGCGCGACGACCTTGAAA</u>	-832 bp relative to <i>fur</i> ORF
P8	<u>TGCGGCCTCGAACTTCGACT</u>	+901 bp relative to <i>fur</i> ORF

^a Underlining indicates common sequences used for PCR; bold indicates unique bar-coding sequences.

1.0) from LS4D modified to contain 0.2% yeast extract, were electroporated with 5 µg of the knockout vector (pMO707) and 1 µg lambda DNA, as an additional substrate for nucleases in the recipient. Prior to electroporation, the cells were washed twice with ice-cold 1 mM MgCl₂-10% (vol/vol) glycerol. Electroporations were carried out in a total volume of 75 µl with a BTX electroporation pulse generator, model ECM630 (Genetronix, San Jose, CA). The parameters obtained for the electroporations were 1.75 kV, 25 µF, and 250 Ω. Immediately following transformation, the cells were recovered in 1 ml LS4D medium supplemented with 0.2% (wt/vol) yeast extract. Following 4 h of incubation at 37°C, the cultures were diluted to 5 ml in the same medium containing 400 µg G418/ml and allowed to grow overnight. The next day the transformation cultures were plated (1 ml/plate) on solidified LS4D plus 0.2% (wt/vol) yeast extract medium containing 400 µg G418/ml. Resulting transformants were analyzed for deletion of the *fur* gene via PCR that targeted genome regions outside of the *fur* knockout cassette (primers P7 and P8 [Table 2]) and Southern analyses of genomic DNAs. One confirmed *fur* deletion mutant was selected and designated JW707.

Nucleic acid procedures. Genomic DNA was extracted with a Wizard genomic purification kit (Promega, Madison, WI). Southern analyses employed Zeta-Probe (Bio-Rad, Hercules, CA) membranes and were performed according to the manufacturer's instructions. For Northern analysis, RNA was isolated from exponential-phase (OD₆₀₀ ~ 0.4) cultures using RNawiz (Ambion, Austin, TX) according to the manufacturer's protocol. Prior to electrophoresis, contaminating DNA was removed from the RNA preparation using DNA-free DNase (Ambion). A total of 10 µg RNA per lane was separated in a gel of 1.2% (wt/vol) agarose with 1× formaldehyde-MOPS-EDTA sodium acetate buffer (Sigma, St. Louis, MO). Following electrophoresis, the RNA was transferred to Zeta-Probe membranes using the downward transfer method described in Ambion technical bulletin 169. The RNA was then permanently affixed to the membrane via UV cross-linking at 120 mJ/cm².

Southern and Northern hybridizations were performed overnight at 42°C using 5 µl ³²P-labeled PCR products in ULTRAhyb solution (Ambion). Blots were washed twice in 2× SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7)-0.1% (wt/vol) SDS for 5 min and twice in 0.1× SSC-0.1% (wt/vol) SDS for 15 min before being placed on film. Probes for both Southern and Northern analyses were generated via PCR (see Table 3 for primer sequences and product sizes) and labeled with [^α-³²P]dCTP using a Prime-It II random primer labeling kit (Stratagene).

Phenotypic analysis. Unless otherwise noted, 2% (vol/vol) subcultures from active cultures in standard Yen45 medium were used as the inoculum for phenotypic analyses. Response to iron limitation was monitored via optical density at 600 nm in Yen45 medium modified by omission of all known sources of iron and addition of 6, 3, or 1 µM FeCl₂. The inocula for testing the growth responses to limiting iron were from medium containing 1 µM added FeCl₂. Modified Yen45 lacking added iron was analyzed by inductively coupled plasma spectrometry by the University of Missouri Agriculture Experiment Station Lab and found to contain 0.3 µM iron. To determine resistance to MnCl₂, CFU were counted following a 112-h exposure to 20, 30, or 40 mM MnCl₂. Growth responses to high concentrations of nitrate and nitrite were monitored via optical density of LS4D cultures amended to contain 50 or 100 mM nitrate or 2 or 5 mM nitrite at time zero. Responses to increased osmolarity were determined via optical density and microscopic observations following the addition of 300 mM NaCl or KCl (46) at the time of subculture. Osmoprotection was assayed by the addition of 2 mM glycine betaine at time zero to subcultures containing 300 mM NaCl or KCl.

Microarray analysis. Volumes of 600 ml of *D. vulgaris* Hildenborough wild type or JW707(Δ*fur*) were grown in LS4D as 100-ml batches in six 125-ml bottles

to a cell density of ca. 3 × 10⁸ cells/ml (OD₆₀₀ ~ 0.4). The 100-ml aliquots were used as inocula for six replicate bottles, each containing 900 ml LS4D, and the cultures were grown to log phase at 30°C in anaerobic chambers. At log phase (OD₆₀₀ = 0.38 for WT and 0.33 for JW707), 250 ml of each culture was harvested for sampling. For iron-limited experiments, the same protocol was followed with modified LS4D containing 5 µM FeCl₂ instead of 60 µM. Cultures were sampled at log phase, OD₆₀₀ = 0.13 for WT and 0.17 for JW707. Cell harvesting, RNA extraction, and microarray analyses were carried out as described previously (46).

RESULTS

Mutagenesis of *D. vulgaris fur* gene. Plasmid pMO707 (Fig. 1) was transferred to wild type *D. vulgaris* via electroporation. Positive (pSC27) and negative (lambda DNA) control DNAs were also electroporated using 10⁹ to 10¹⁰ recipient cells for each transformation. To select for transformants, cells were plated on LS4D medium modified to contain 0.2% (wt/vol) yeast extract and 400 µg G418/ml and colonies appeared after four days at 30°C. The transformation efficiency for pMO707 was ca. 1.2 × 10⁻⁷ per recipient cell, while the efficiency for the

TABLE 3. PCR primers for gene probe construction

Target	Primer	Sequence (5'-3')	Product size (bp)
DVU0942 (<i>fur</i>)	fur-65F	AGCGGATGCTCATCGT CGAT	334
	fur-399R	GAGATACATCCGGTGC GAGG	
DVU2680 (<i>fld</i>)	fld-16F	ATCGTCTACGGTTCCA CCAC	294
	fld-310R	CCCCGCAGAAGTACTC GTAG	
DVU2410 (<i>sodB</i>)	sodB-88F	GCCTACTACGCCAACC TCAA	400
	sodB-488R	CATACGTCGATGGTCA GCAC	
DVU2572 (<i>feoA</i>)	feoA1-13F	ATCTCCCTGCGGCA GTTG	230
	feoA1-243R	GTCGGCTTCCATCACT TCAA	
Km ^r from pSC27	Km-intF	AGATCTGATCAAGAGA CAGGATGAG	729
	Km-intR	CTCTTCAGCAATATCA CGGGTAG	




FIG. 1. Map of pMO707 containing the *fur* deletion construct. A 2,706-bp PCR product containing the pSC27 Km^r determinant flanked by DNA sequences upstream (768 bp) and downstream (831 bp) of the *D. vulgaris fur* gene was inserted into the EcoRV site of the pBluescript SK(+) multicloning site. Both pBluescript and pMO707 are unstable in *D. vulgaris*. Numbers indicate positions of sequences upstream and downstream of the *fur* gene; BC indicates molecular barcodes allowing mutant tracking in a mixed population.

stable plasmid pSC27 was 1.4×10^{-6} per recipient cell. No G418 resistant colonies resulted from plating the negative control transformation. Colonies from the pMO707 transformation were subcultured into liquid medium of the same composition as that used for plating and a second single-colony isolation was made. Deletion of the *fur* gene via marker exchange was verified by a change in the size of a PCR product generated from primers complementary to regions up- and downstream of the *fur* gene (Fig. 2A). While the resulting PCR product from the wild-type *fur* region was 2168 bp, the product from two selected transformants (JW706 and JW707) was 671 bp larger (2,839 bp total). This difference correlates to the

larger size of the Km^r determinant replacing the *fur* gene. Southern analysis with probes internal to *fur* or the Km^r determinant also corroborated replacement of the *fur* gene with the Km^r determinant (data not shown) and JW707 was selected for further analysis.

Northern analysis. RNAs from exponential-phase ($OD_{600} \sim 0.4$) wild-type and JW707 cultures grown in standard LS4D medium were analyzed for the expression of the *fur* gene as well as other genes predicted to be regulated by iron (61): *fld* (DVU2680; flavodoxin), *sodB* (DVU2410; superoxide dismutase), and *feoA* (DVU2572; ferrous iron transporter) (Fig. 2B). Hybridization analysis with the 334-bp *fur* probe indicated the presence of a single ~ 500 -bp transcript for the wild type, but as expected, no signal was present in the JW707 lane (Fig. 2B, lane 2). This transcript size correlates with the predicted *fur* ORF, including a possible promoter sequence. For the *fld* gene, a weak band corresponding to a ~ 650 -nucleotide transcript was observed for the wild type, while an intense band was present for JW707. While the *fld* ORF was annotated to be 447 bp, the Fur binding site was predicted to be 182 bp upstream (61). This hybridization signal is indicative of a sharp increase in expression. No transcript was evident for the *sodB* gene in either the wild-type or the JW707 samples, indicating expression at a level below detection (Fig. 2B). Results from the *feoA* probe indicated a large signal smear for JW707, with limited hybridization for the wild type. This lack of hybridization specificity may be due to the operon of which *feoA* is a part (DVU2571 to DVU2572), as well as the two genome copies of the *feoA* gene (DVU2572 and DVU2574). Predicted operon arrangements for the targeted genes are indicated in Table 4.

Response to iron. Growth curves for both the wild type and JW707 in Yen45 ($30 \mu\text{M FeCl}_2$) were identical, indicating that the *fur* gene is not essential for anaerobic dissimilatory sulfate reduction by *D. vulgaris* under iron-replete conditions (Fig. 3A). To test the effect of iron limitation on the *fur* mutant, growth was analyzed in Yen45 medium containing 1 and $6 \mu\text{M}$ added FeCl_2 . Inocula used in this experiment were 2% (vol/vol) of active cultures in medium containing $1 \mu\text{M}$ added FeCl_2 to limit carryover of excess iron. Resulting growth curves for




FIG. 2. (A) PCR analysis of Km^r *D. vulgaris* transformants using primers targeting regions outside of the *fur* knockout cassette (P7/P8). Lane 1, wild type; lane 2, transformant A; lane 3, transformant B; lane 4, pMO707; lane 5, no DNA; lane M, 1-kb marker. An increase in product size from 2,168 bp to 2,839 bp indicates exchange of the *fur* gene for the Km^r determinant, creating JW707. (B) Northern analyses of total RNA (10 μg) from JW707. Lane 1, wild type; lane 2, JW707; lane M, RNA marker. Probes used for hybridization are indicated above the blot. nt, nucleotides.

TABLE 4. Effect of Δfur on expression of selected genes in *D. vulgaris* Hildenborough^a

Category and ORF ^b	Gene ^c	Annotated function	Expression ratio for comparison (strain, FeCl ₂ concn [μ M]) ^d				
			Fur, 60/WT, 60	Fur, 5/WT, 5	WT, 5/WT, 60	Fur, 5/Fur, 60	
Genes differentially expressed in the Δfur strain regardless of iron concn ^e							
↓	DVU0273*	NA	Conserved hypothetical protein	4.6	1.7	1.8	-0.6
↑	DVU2158	NA	Hypothetical protein, phage #5	2.5	1.9	0.4	0.3
↓	DVU2378*	<i>foxR</i>	Transcriptional regulator, AraC family	3.3	2.8	1.2	1.2
	DVU2379	<i>pqqL</i>	Peptidase, M16 family, putative	2.5	2.1	0.8	0.8
	DVU2380	<i>aipX</i>	ABC transporter, ATP-binding protein	3.8	2.5	1.1	0.4
	DVU2381	NA	Conserved hypothetical protein	4.4	3.4	1.1	0.7
↓	DVU2382	NA	Conserved domain protein	3.4	2.4	1.0	0.5
	DVU2383	NA	<i>tonB</i> -dependent receptor domain protein	5.2	3.3	1.2	ND
	DVU2384	COG- <i>oppA</i>	ABC transporter, periplasmic substrate-binding protein	2.2	2.3	0.4	0.9
	DVU2385	NA	ABC transporter, permease protein	1.1	1.1	1.6	1.6
	DVU2386	<i>oppC</i>	ABC transporter, permease protein	0.7	1.0	0.1	0.6
	DVU2387	NA	ABC transporter, ATP-binding protein	1.2	1.4	0.6	0.9
	DVU2388	<i>tolQ1</i>	TolQ protein	2.8	2.1	0.8	0.5
	DVU2389	<i>tolR</i>	Biopolymer transport protein, ExbD/TolR family	2.2	1.5	0.5	0.1
↓	DVU2390	COG- <i>tonB</i>	TonB domain protein	2.3	1.2	0.7	-0.1
↑	DVU2571	<i>feoB</i>	Ferrous iron transport protein B	4.6	1.2	3.0	-0.1
	DVU2572	<i>feoA</i>	Ferrous iron transport protein A	4.1	1.7	2.2	0.4
↑	DVU2573	NA	Hypothetical protein	3.1	1.7	1.4	0.2
↑	DVU2574*	<i>feoA</i>	Ferrous iron transporter component FeoA	2.4	1.4	0.4	-0.2
↓	DVU2680*	<i>fld</i>	Flavodoxin, iron repressed	No data	2.0	2.8	No data
↑	DVU2681	NA	Hypothetical protein	5.3	2.3	2.7	ND
Additional genes predicted to be part of the Fur regulon of <i>D. vulgaris</i> Hildenborough ^f							
↓	DVU0763*	<i>gdp</i>	GGDEF domain protein	3.9	1.1	1.3	-1.0
↓	DVU0303	<i>genZ</i>	Hypothetical protein, GenZ	1.3	-0.7	2.2	0.1
↓	DVU0304*	<i>genY</i>	Hypothetical protein, GenY	-0.8	-0.7	-0.2	-0.2
↓	DVU3123	<i>hdd</i>	HD domain protein	1.3	0.2	0.4	-0.4
↓	DVU3124	NA	Hypothetical protein	2.2	0.1	0.6	-1.2
↓	DVU3330*	NA	Hypothetical iron-regulated P-type ATPase	1.9	0.3	2.0	0.5
↓	DVU3331	NA	Hypothetical protein	2.4	0.2	2.2	0.1
↓	DVU3332	NA	Heavy metal-translocating P-type ATPase	1.2	ND	1.0	-0.2
↓	DVU3333	NA	Hypothetical protein	1.8	0.2	1.8	0.3

^a Underlined entries (ORFs in the first column and corresponding ratios) do not meet the significance criteria (\log_2 ratio ≥ 1.6) selected for this study but were predicted to be in the same operon as genes that do or genes in the Fur regulon.

^b Arrows indicate the predicted direction of transcription and operon arrangement. Asterisks indicate ORFs predicted to be preceded by a putative δ -proteobacterial Fur binding site (61).

^c NA, not annotated. COG, clusters of orthologous groups.

^d \log_2 ratios of transcriptional responses. Only ratios of ≥ 1.6 are considered significant, although operon arrangement is noteworthy. ND, no transcript differences were detected between the two conditions.

^e Boldface indicates differential expression consistent with genes regulated by Fur regardless of iron concentration.

^f Boldface indicates differential expression consistent with genes regulated by iron-bound Fur.

both iron levels were similar between the wild type and JW707 (data not shown). The Fur mutant did not exhibit a growth advantage over the wild type when iron was limiting.

Metal sensitivity. Since a common trait of bacterial Δfur mutants is increased resistance to Mn(II) (8, 30, 32, 40, 55), the effect of increasing concentrations of MnCl₂ on the growth of both wild-type and JW707 strains was tested. Excessive precipitation caused by the addition of MnCl₂ to the growth medium (data not shown) mandated that CFU be monitored as an indicator of Mn(II) effects on growth. Following 112 h of

exposure to 20, 30, or 40 mM MnCl₂, cultures were plated on Yen45 medium. Wild-type untreated cultures had more CFU than untreated JW707; however, with 40 mM MnCl₂, the number of wild-type CFU declined from 1.3×10^8 to 1.4×10^5 (Fig. 4). By comparison, the numbers of CFU for JW707 achieved in cultures exposed to MnCl₂ were similar, with 3.7×10^7 CFU in the untreated control and 1.8×10^7 CFU after exposure to 40 mM MnCl₂ (Fig. 4).

Response to osmolarity shock. Previous studies with wild-type *D. vulgaris* indicated that salt stress reduced growth rate




FIG. 3. Growth curves of the *D. vulgaris* wild type (open symbols) and mutant strain JW707 (filled symbols) under various conditions. (A) Growth under iron-replete conditions ($30 \mu\text{M FeCl}_2$). (B) Response to osmolarity stress, with sodium as 300 mM NaCl (circles) or 300 mM NaCl plus 2 mM glycine betaine (squares). (C) Response to osmolarity stress, with potassium as 300 mM KCl (circles) or 300 mM KCl plus 2 mM glycine betaine (squares). (D) Response to nitrite at 2 mM NaNO_2 (circles) and 5 mM NaNO_2 (squares). Curves are representative of three trials at 37°C .

and increased cell length and that these effects could be countered by addition of the osmoprotectant glycine betaine (46). Interestingly, the putative Fur regulon members were among the most highly differentially transcribed genes during stress with NaCl or KCl (46). Therefore, the response of the Δfur mutant to salt stress was tested. Growth curves of cultures in medium supplemented with 300 mM additional NaCl or KCl (Fig. 3B and C, respectively) were similar and growth rates decreased for both the wild type and JW707. Microscopic observations of the salt-stressed cultures showed that $91.4\% \pm 2.8\%$ (standard deviation) of wild-type cells were elongated from $1.5 \mu\text{m}$ to $5 \mu\text{m}$ or more. Although JW707 cells were not counted, the fraction elongated appeared to be similar to that of the wild type. Addition of 2 mM glycine betaine with the salt protected the wild type from the decrease in growth rate (Fig. 3B and C) and restored the cell morphology to normal. However, the growth rate of JW707 remained 75% of the rate of uninhibited cells regardless of the presence of 2 mM glycine betaine. Interestingly, microscopic observations indicated that

the osmolyte apparently restored the mutant cell lengths to $\sim 1.5 \mu\text{m}$ (data not shown).

Response to nitrate/nitrite. Tests of JW707 growth responses to nitrate or nitrite stress were prompted by data from proteomic and microarray analyses of wild-type *D. vulgaris* (35, 59). The transcript analyses indicated upregulation of both the *fur* regulon and genes predicted to be involved in iron binding, particularly under nitrite stress. Significant differences in growth between the wild type and JW707 were not observed in Yen45 medium containing 50 mM NaNO_3 (data not shown). Increasing this NaNO_3 concentration to 100 mM did not have an effect on the final growth extent, but a 60-h lag phase for the wild type and an 80-h lag phase for JW707 did occur (data not shown). Growth of the wild type in Yen45 medium supplemented with 2 mM NaNO_2 resulted in a lag phase of 60 h, whereas no growth was detected in medium supplemented with 5 mM NaNO_2 after 117 h (Fig. 3D). In contrast, both 2 and 5 mM NaNO_2 completely inhibited the growth of JW707. Since comparable cell numbers were used as inocula for the parent and mutant, the results indicate an increased sensitivity to nitrite by the Δfur mutant.

Expression profile of JW707. Transcriptional arrays covering 3,482 of the 3,531 (98.6%) protein coding sequences in the *D. vulgaris* genome were used to identify genes affected by deletion of the *fur* gene. In addition, the effect of iron concentration on the global transcription of the *fur* mutant was also determined by performing two separate experiments with growth medium containing $60 \mu\text{M}$ and $5 \mu\text{M}$ added FeCl_2 . Samples were taken at similar optical densities for both JW707 and the parental wild-type strain, with differential gene expression calculated as $\log_2(\text{transcripts of JW707}) - \log_2(\text{transcripts of wild type})$. Following normalizations for signal intensities (13) and sector-




FIG. 4. Growth response of the *D. vulgaris* wild type (open symbols) and mutant strain JW707 (filled symbols) to MnCl_2 treatment. Curves are representative of three trials at 37°C .




FIG. 5. (A) Genes differentially expressed threefold in the Δfur mutant, JW707, compared to wild-type cells, under iron-replete (+Fe; 60 μM) and iron-limited (-Fe; 5 μM) conditions. (B) Overlap of genes differentially expressed threefold in JW707 iron-limited (-Fe; 5 μM) conditions compared with JW707 iron-replete conditions versus those differentially expressed in iron-limited wild-type cells compared to iron-replete wild-type cells.

based artifacts, the significance of the ratios was calculated as a Z score. Generally, ratios of ≥ 1.6 (≥ 3 -fold change in expression) were selected for further analysis.

Evidence that Fur is a global regulator in *D. vulgaris* derives from the observation that changes in gene expression with *fur* deleted were identified in 12 functional categories based on the annotation by The Institute for Genomic Research. Transcript analysis revealed 34 and 50 genes differentially expressed at least threefold in response to the *fur* deletion (compared to the wild type) under iron-replete (60 μM) and iron-limited (5 μM) conditions, respectively (Fig. 5A). Comparison of the two data sets indicated that expression levels for 13 genes were affected under both iron conditions (Table 4). Under iron-replete conditions, 30 genes were up-regulated and 4 genes were down-regulated, with 44% of the total population predicted to encode hypothetical or conserved hypothetical proteins. Under iron-limiting conditions, 32 genes were up-regulated and 18 genes were down-regulated, with 30% of the total population predicted to encode hypothetical or conserved hypothetical proteins.

When JW707 responses to limiting iron were compared to its responses in iron-replete conditions, 22 genes were increased in expression, compared to 22 that were decreased. In contrast, when the wild type was subjected to limited iron, 50 genes were increased in transcription and only 18 were decreased relative to iron-replete cells (Fig. 5B). Thus, the genes differentially expressed in JW707 may be candidates for iron regulation. Genes expressed differently in the wild type should include those regulated by Fur and those regulated by iron concentration. Curiously, only three genes meeting the stringent criterion of a threefold increase in expression were common to iron-restricted JW707 and iron-restricted wild type: these were annotated as rubrerythrin (*rbr2*; DVU2318), chemotaxis protein (*cheY2*; DVU2073) (Table 5, category IV), and an ABC transporter permease protein (DVU2385) (Table 4).

Transcriptional profile of the predicted Fur regulon. Computational analysis of the upstream regions of *feoAB* operons from multiple δ -proteobacterial genomes revealed a conserved 17-bp palindromic motif corresponding to a putative Fur binding site (61). A genomic scan of the *D. vulgaris* genome with this motif revealed 17 sites upstream of eight genes or operons, constituting the putative Fur regulon (61). The prediction that these sites are operator sites for Fe(II)-bound Fur appears to

have been supported by some of the transcriptional responses of the wild type compared to those of the Fur mutant (Table 4). Comparison of the transcripts from the wild type under iron-limited versus iron-replete conditions also supported the prediction that Fur is not autogenously regulated in *D. vulgaris* (61) (data not shown). The lack of correlation between expression changes in the Fur regulon and the *fur* gene in microarray data from stressed *D. vulgaris* cells (35, 46) also suggests that *fur* is not autogenously controlled.

The DVU2681 gene, encoding a hypothetical 60-amino-acid protein, exhibited the most up-regulation in the transcriptional profiling of JW707: \log_2 ratios of 5.31 and 2.30 under iron-replete and -restricted conditions, respectively. It should be noted that DVU2681 is located directly downstream of the flavodoxin gene (DVU2680) that is annotated as "iron repressed" and, by Northern analysis, shown to be greatly increased in transcription in the Δfur mutant (Fig. 2B). However, DVU2681 is transcribed in the opposite orientation. Other putative Fur-regulated operons, such as those encoding the annotated GGDEF domain protein (DVU0763) and the HD domain protein (DVU3123) (Table 4), did not appear to respond to decreased iron concentration. In the putative *genYZ* operon, only *genZ* (DVU0303) expression was consistent with iron-bound Fur repression. The latter two genes were predicted to be members of a Fur regulon unique to metal-reducing δ -proteobacteria, though possible functions are unknown (61). In contrast, the hypothetical iron-regulated P-type ATPase gene (DVU3330) that was predicted to be monocistronic responds as if it were part of an operon of two to four genes (Table 4). A similar expansion of Fur or iron influence on expression can be seen downstream of the *foxR* regulatory gene (DVU2378), which appears to include 12 genes (Table 4). This cluster is composed of ABC transporters (DVU2380, DVU2384 to DVU2387), the biopolymer transporters (ToIRQ; DVU2388 to DVU2389), and an energy-generating mechanism for transporting polymers across the membrane (TonB [DVU2390] and a TonB receptor [DVU2383]). While these genes have been shown to be involved in iron acquisition via siderophore production and transport in many bacteria (39, 44, 56), *D. vulgaris* has not been shown to produce siderophores (unpublished data), nor have genes for siderophore biosynthesis been recognized in the genome. Whether the transcriptional changes are a direct response to Fur deletion or are mediated by influences on activities of other regulators, such as FoxR, will need to be addressed with further experimentation.

Other iron-related genes. Transcriptional responses of genes that might be predicted to be involved with iron metabolism from the reported roles of their orthologs in other bacteria were examined (data not shown). Expression profiles of a putative siderophore uptake system encoded by DVU0650 to DVU0646 and of two iron storage proteins, bacterioferritin (DVU1397) and ferritin (DVU1568), did not support a role for Fur regulation or a clear response to iron concentrations. Two proteins requiring iron for function, Fe hydrogenase (DVU1771) and ferredoxin II (DVU0305), were also not significantly altered in expression in the absence of Fur.

Selected genes believed to be involved in the oxidative stress response of *D. vulgaris* were also examined for transcriptional responses in the deletion mutant (data not shown). Only the relative transcription for the gene annotated to encode alkyl

TABLE 5. *D. vulgaris* Hildenborough transcriptional responses to Δfur and FeCl₂ concentrations suggesting different regulatory systems

Category and ORF ^a	Gene ^b	Annotated function	Expression ratio for comparison (strain, FeCl ₂ concn [μ M]) ^c				
			Fur, 60/WT, 60	Fur, 5/WT, 5	WT, 5/WT, 60	Fur, 5/Fur, 60	
I. Repressed by iron-bound Fur^d							
↑	DVU3300	NA	Hypothetical protein	2.7	No data	3.8	-0.3
↑	DVU3301	NA	Hypothetical protein	2.0	-1.2	3.5	0.2
II. Repressed by iron-free Fur							
↑	DVU0459	NA	Hypothetical protein	ND	1.6	-1.3	0.5
	DVU0465	<i>trpE</i>	Anthranilate synthase, component 1	ND	2.5	-1.2	1.6
	DVU0466	<i>trpG</i>	Anthranilate synthase, glutamine amidotransferase component	ND	1.9	-1.1	1.0
	DVU0467	<i>trpD</i>	Anthranilate phosphoribosyltransferase	-0.1	1.8	-1.2	0.8
	DVU0468	<i>trpC</i>	Indole-3-glycerol phosphate synthase	-0.1	1.7	-1.1	0.9
	DVU0469	<i>trpF1</i>	<i>N</i> -(5-Phosphoribosyl)anthranilate isomerase	0.4	2.0	-1.3	0.6
	DVU0470	<i>trpB2</i>	Tryptophan synthase, beta subunit	0.3	2.0	-1.5	0.5
	DVU0471	<i>trpA</i>	Tryptophan synthase, alpha subunit	-0.2	1.6	-1.2	0.8
↓	DVU2449	<i>metK</i>	<i>S</i> -Adenosylmethionine synthase	-0.2	1.8	-1.6	0.5
III. Induced by iron-free Fur							
↓	DVU0241	NA	MTH1175-like domain family protein	ND	-1.8	1.6	0.3
↓	DVU0593	<i>lysE</i>	L-Lysine exporter, putative	0.1	-1.8	2.2	ND
↓	DVU0811	<i>dnaK</i>	DnaK protein	-0.5	-1.7	2.0	0.6
↓	DVU1401	NA	Membrane protein, putative	0.2	-2.1	2.8	0.1
↓	DVU1444	<i>flgD</i>	Basal-body rod modification protein FlgD	1.1	-2.3	3.3	-0.3
↓	DVU2886	NA	Transcriptional regulator, AraC family	-0.2	-2.2	2.9	0.5
IV. Iron repressed							
↓	DVU2073	<i>cheY-2</i>	Chemotaxis protein CheY	0.2	0.2	1.6	1.7
↓	DVU2318	<i>rbr-2</i>	Rubrythrin, putative	ND	0.4	2.2	2.5
↑	DVU2441	<i>hspC</i>	Heat shock protein, Hsp20 family	-0.2	-1.1	1.9	0.8
↑	DVU2442	NA	Heat shock protein, Hsp20 family	-0.1	-1.0	2.2	1.1
↓	DVU2494	NA	Peptidase, M48 family	0.1	-0.5	2.0	1.3
V. Iron induced							
↓	DVU0186	NA	Conserved hypothetical protein	0.5	0.6	-2.2	-2.0
↓	DVU0607	<i>ahcY</i>	Adenosylhomocysteine	-0.1	-0.2	-1.6	-1.8
↓	DVU0997	<i>metF</i>	5,10-Methylenetetrahydrofolate reductase	0.2	0.1	-1.6	-1.8
↓	DVU1257	NA	RNA-binding protein	0.7	0.5	-1.4	-1.5
↓	DVU3371	<i>metE</i>	5-Methyltetrahydropteroyltriglutamate-homocysteine <i>S</i> -methyltransferase	0.2	-0.5	-2.2	-2.6
↓	DVUA0019	NA	Type II DNA modification methyltransferase, putative	-0.4	-0.3	-2.3	-2.1
↓	DVUA0067	NA	Membrane protein, putative	0.5	-0.7	-2.3	-3.2
↓	DVUA0068	NA	Membrane protein, putative	0.3	-0.7	-1.9	-2.9
↓	DVUA0069	NA	GtrA family protein, selenocysteine containing	0.3	0.5	-2.1	-1.7

^a Arrows indicate direction of transcription of operons or genes.

^b NA, not annotated.

^c Log₂ ratios of transcriptional responses. Only ratios of ≥ 1.6 are considered significant, although operon patterns are noteworthy. Boldface shows the number used as the primary criterion for grouping. ND, no transcript differences were detected between the two conditions.

^d Not predicted by original motif searches (62).

hydroperoxide reductase C (DVU2247) was consistent with iron-bound Fur-dependent repression, although the changes did not meet our cutoff for significance. This gene product is reported to reduce hydrogen peroxide and to protect the cell from reactive oxygen species formed while iron acquisition systems are induced (inactive Fur repressor). Changes in the expression of genes for the putative cytochrome *d* ubiquinol oxidase (DVU3270 to DVU3271), superoxide dismutase (DVU2410), and catalase (DVUA0091) were not consistent with classical Fur-mediated regulation. As has been shown in other systems (4, 11, 25), the catalase gene transcription was

increased by limiting iron, an example of regulation of a gene located on the megaplasmid in *D. vulgaris*.

Other potential modes of maintaining iron homeostasis. Iron-responsive regulation through the Fe(II)-bound Fur repressor is the most commonly identified microbial mechanism for maintaining iron homeostasis (31). However, a second Fe(II)-responsive repressor, DtxR, has been identified, primarily in gram-positive bacteria (31), and a role for Fe(III)-specific two-component regulatory systems has been recognized (56). From sequence analysis of *D. vulgaris*, homology to DtxR proteins has not been recognized, nor have Fe(III)-specific histi-

dine kinases and response regulators been annotated. However, the expression of a number of genes would be consistent with alternative modes of regulation (Table 5), as have been identified recently for *Helicobacter pylori* (14, 21). Category I illustrates the predicted pattern for Fe(II)-bound Fur regulation: increased expression in the absence of Fur in plentiful iron and when Fe(II) is limiting in wild-type cells. Other candidates are shown in Table 4. The two hypothetical genes shown in category I (Table 5) are highly regulated, but possible functions remain obscure. Category II is exemplified by amino acid biosynthesis genes for tryptophan and for methionine. These genes are up-regulated in the absence of Fur and low iron and decreased in expression in the presence of Fur and low iron, a pattern consistent with repression by iron-free Fur. Category III lists a few examples of genes that exhibit expression patterns that could be compatible with iron-free Fur induction. In limiting iron, low levels of expression are observed in the absence of Fur, but high levels of expression are measured when Fur is present. In wild-type cells with 5 μM FeCl_2 , Fur would induce but the addition of 60 μM FeCl_2 would prevent induction. The apparent positive regulation by iron-free Fur in *D. vulgaris* requires confirmation before mechanisms are sought.

Regulation in response to iron concentrations that is not mediated by Fur may also be apparent in *D. vulgaris*. Categories IV and V show transcription patterns that seem consistent with iron repression and iron induction, respectively. For iron repression, candidate regulated genes were increased in expression in limiting iron regardless of the presence of Fur. For iron induction, the opposite pattern of transcript levels was seen. The two genes whose expression was found to be increased over threefold in both iron-limited and iron-replete conditions, *rbr2* and DVU2541, are among those in category IV (Table 5). Curiously, category V includes several genes present on the 200-kb megaplasmid in wild-type *D. vulgaris*.

DISCUSSION

Genetic system. Here we describe a variation on the genetic system for *D. vulgaris* (26). We constructed the knockout cassette by fusion PCR to delete target genes via double recombination following electroporation. The development of this gene deletion method for *Desulfovibrio* was necessitated by a gene reversion event observed following plasmid interruption of the *cycA* gene in *D. desulfuricans* G20 (57). Due to the possibility of verifying deletion mutants within 2 weeks following cassette construction, this method can be used for rapid production of *D. vulgaris* deletion mutants. In 1997, Fu and Voordouw described the first case of targeted gene deletion in *D. vulgaris* (26). This method is dependent on Cm^r selection and SacB counterselection following two separate recombination events and has been successful in generating at least 11 *D. vulgaris* deletion mutants (reference 5 and references therein; 75). Because the SacB counterselection procedure involves DNA transfer by conjugation as well as lengthy transformant screening, its use in high-throughput mutant construction is not practical.

A clean selection method is also key for the success of a genetic system, and developing such a method for *Desulfovibrio* has been problematic due to broad-range antibiotic resistance.

While the method described by Fu and Voordouw utilizes the Cm^r determinant (26), chloramphenicol resistance has been problematic in our hands. We have found that 400 μg of G418 (gentamicin) per ml of LS4D medium provides clean selection when the Km^r determinant is used. The utility of this method is illustrated by the 1.25×10^{-7} transformation efficiency obtained and the successful deletion of the *fur* gene in all three of the transformants screened via phenotypic (MnCl_2 resistance) and molecular (PCR and Southern analysis) methods.

The method described here is also dependent on DNA delivery by electroporation, which had not been successful in *D. vulgaris* until now. We added a high concentration of deletion vector as well as lambda DNA to an electroporation procedure first described for *Desulfovibrio fructosivorans* (64). The intrinsic nucleases of *D. vulgaris* likely linearize the plasmid and promote double recombination events and thus marker exchange between the vector and genome (5). Other deletion mutants screened in our laboratory using this deletion procedure follow the same pattern (data not shown).

Physiology of JW707. The similar decreases in growth rate for both the wild type and JW707 with 5 μM added FeCl_2 indicated that both cultures were limited for iron, but not differentially (data not shown). Thus, regardless of iron concentration, growth of the Δfur mutant was neither inhibited nor promoted by the derepression of putative ferrous iron uptake genes (*feoAB*) (Fig. 2B and Table 4). Similarly, neither *Dichelobacter nodosus* nor *Shewanella oneidensis* Fur mutants exhibited a detectable phenotype when grown anaerobically (52, 71). Without oxygen present, toxic radical formation resulting from increased ferrous iron and the Fenton reaction would be less likely. However, further tests are needed to determine if the cellular iron content of the *D. vulgaris* Fur mutant differs from that of the wild type.

Osmotic stress. Salt stress in *Bacillus subtilis* has been shown to induce iron acquisition systems; therefore, salinity has been proposed to cause iron limitation (37, 70). Analysis of the *D. vulgaris* transcriptome and proteome under both NaCl and KCl stress indicated the same phenomenon, Fur regulon induction in response to salt stress (46). In contrast, physiological studies indicated growth inhibition of JW707 equivalent to that of the wild type upon exposure to 300 mM NaCl or KCl (Fig. 3). Therefore, increased expression of the Fur regulon is not sufficient for overcoming this osmotic stress in *D. vulgaris*.

$\text{NO}_3^-/\text{NO}_2^-$ stress. The effect of nitrate on *D. vulgaris* growth is of specific concern for environmental bioremediation applications. High levels of nitrate in uranium-contaminated have been documented sites (<http://www.esd.ornl.gov/nabirfrc/>) (20), and nitrate is believed to have an inhibitory affect on sulfate reduction through the intermediate nitrite (42, 49, 50). A previous proteomic analysis of wild-type *D. vulgaris* showed that the presence of 105 mM NaNO_3 induced proteins involved in the ionic stress response (59), although transcriptional profiling has indicated a response unique to nitrate that is not a composite of salt and nitrite (unpublished data).

The increased sensitivity to nitrite stress for JW707 was unexpected based on previous transcript analyses of wild-type *D. vulgaris* (35). An increased expression of the Fur regulon in the wild type was observed under nitrite stress (35). Therefore, derepression of the regulon through deletion of the repressor was expected to provide an advantage for JW707 exposed to

nitrite. However, the Fur mutant was completely inhibited by 5 mM NaNO₂ (Fig. 3D). Like JW707, an *Escherichia coli* Δfur mutant was also more sensitive to NaNO₂ regardless of the upregulation of genes involved in iron uptake (48). This increased expression of the Fur regulon following 1 mM NaNO₂ treatment was hypothesized to be a consequence of NO nitrosylation of the Fe(II) in Fur that inactivated the repressor and likely other iron-containing genes (15). Direct evidence for this mechanism of iron modification in nitrite-treated *D. vulgaris* remains to be obtained.

Transcriptional profile of JW707. While the overall physiology of *D. vulgaris* was not dramatically affected by deleting the *fur* gene, a diverse transcriptional response occurred for JW707 compared to the wild type. In many bacteria, Fur negatively regulates siderophore production and transport (reference 67 and references therein). Despite the up-regulation of a 12-gene cluster containing genes predicted to be involved in siderophore uptake (e.g., TolRQ) (Table 4), no genes for siderophore production have been annotated in the *D. vulgaris* genome. This may allow *D. vulgaris* to save energy by simply stealing iron-complexed siderophores produced by other bacteria, as suggested for the spirochete *Leptospira biflexa* (43). In other bacteria, similar TonB/ABC and biopolymer uptake systems have also been shown to transport vitamin B₁₂, phages, colicins, and maltodextrins into the cell (51, 54). Further studies are needed to determine what role the large (12-gene) cluster plays in *D. vulgaris*.

The high level of differential expression for the *feoAB* operon, apparently regulated by Fur and its corepressor Fe(II) (Fig. 2 and Table 4), is interpreted as evidence that the FeoAB system is the primary iron uptake mechanism in *D. vulgaris*. This follows from the prediction that Fe(II) predominates in anaerobic environments. However, genes predicted to be involved in iron storage, such as *bfr* and *ftn*, did not exhibit a strong transcriptional response to either iron concentration or the deletion of the *fur* gene.

Oxidative stress. The Fur regulator has been shown to be intricately involved in the oxidative stress response of some bacteria (2, 33, 34). Thus, it can be argued that regulation by Fur is linked to an increased uptake of iron that can generate reactive oxygen species upon oxygen exposure. Despite the anaerobic lifestyle of *D. vulgaris*, it is known to survive exposure to oxygen and contains several oxidative stress response genes within its genome. As such, it was not surprising to find that *ahpC*, *rhl*, *cydA*, and *kata* were differentially expressed at low levels in this study. However, only *ahpC* appeared to be classically regulated by iron-bound Fur (data not shown). The diverse regulatory pattern of the other oxidative stress response genes may be explained by coregulation by PerR, a homolog to Fur that also responds to iron concentration (9, 11). In fact, *ahpC* and *rhl* are predicted to possess a PerR binding site within their promoter regions (61). Coregulation of *kata* and *ahpC* by Fur and PerR has also been proposed in the microaerophilic *Campylobacter jejuni* (3, 38, 73). Further studies are needed to determine if Fur and PerR regulation overlap in *D. vulgaris*.

Other mechanisms of Fur regulation. In addition to the traditional repressor role of iron-bound Fur, alternative activation and repression by iron-free Fur were inferred from the transcription patterns seen in *D. vulgaris*. These two forms of

regulation have recently been described in *Helicobacter pylori* (14, 18, 22). While iron-bound Fur has been shown in the literature to be a positive regulator, it is possible that some of the affected genes are actually repressed by iron-free Fur. This form of Fur regulation was described for the *pfr* (which encodes a ferritin) and *sodB* genes of *H. pylori* (18, 22). Some of the genes for which iron-free Fur repression is consistent in *D. vulgaris* are the seven genes of the tryptophan operon and *metK* (Table 5).

Alternatively, the transcription of flagellar genes (DVU1443 to DVU1445) appeared to be consistent with induction by iron-free Fur in *D. vulgaris*, and DVU1444 expression changes met the criteria used as an example in this category (Table 5). This form of positive regulation by iron-free Fur has been reported only for *flaB*, a major flagellin gene, in *H. pylori* (14). It is tempting to infer that motility may be decreased in the presence of plentiful iron for both *D. vulgaris* and *H. pylori*. Another example of positive regulation by Fur is activation by iron-bound Fur, suggested for a collection of *Neisseria meningitidis* genes involved in both aerobic and anaerobic respiration (17). However, expression patterns suggesting this form of regulation were not observed in this study.

Iron regulation independent of Fur. Differential expression of genes that is independent of Fur implies that other regulators may respond to iron concentrations within the cell. Possible examples of these regulators include PerR, DtxR, NikR, Irr, and RirA (6, 9, 10, 14, 16, 28). However, only *perR* has been annotated in the *D. vulgaris* genome (61), with DVU2318 (encoding a rubrerythrin) being the only gene in the iron-responsive category predicted to be PerR regulated (Table 5) (61). Thus, it appears that other iron-responsive regulators may be present in *D. vulgaris* but have not yet been identified. Based on the iron-dependent regulation of megaplasmid-borne genes (Table 5), these iron-responsive regulators may not be limited to the *D. vulgaris* chromosome but may also be present on the megaplasmid. However, preliminary searches for regulatory motifs upstream of genes showing similar differential expression patterns have been unsuccessful. This is not too surprising because of the small number of candidate genes involved in each category, although both the genes from *D. vulgaris* and the orthologs from *D. desulfuricans* G20 were included in the search (data not shown).

In summary, both the physiological and microarray data indicate a global regulatory role for Fur in *D. vulgaris*, including involvement in the osmotic and nitrite stress responses. While a traditional repressor role for iron-bound Fur was observed, expression data also suggest activation by iron-free Fur. The patterns of apparent transcription regulation presented here require further confirmation but indicate that alternate models of iron homeostasis may be functioning in *D. vulgaris* and possibly anaerobes in general. Specifically, studies to determine the roles of iron-free Fur and the TonB/TolQR system in addition to identification of other iron-dependent regulators are needed. These issues are especially pertinent because of the differential expression of the *D. vulgaris* Fur regulon upon exposure of the cells to various environmental stressors (35, 46, 74). Information from these studies may prove integral in the design of future bioremediation strategies for the removal of heavy metals.

ACKNOWLEDGMENTS

This work was part of the Virtual Institute for Microbial Stress and Survival (<http://vimss.lbl.gov>), supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomics Program: GTL through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy.

We thank Barbara Giles for sharing her technical expertise regarding *Desulfovibrio* mutant construction.

REFERENCES

- Andrews, S. C. 1998. Iron storage in bacteria. *Adv. Microb. Physiol.* **40**:281–351.
- Andrews, S. C., A. K. Robinson, and F. Rodriguez-Quinones. 2003. Bacterial iron homeostasis. *FEMS Microbiol. Lett.* **27**:215–237.
- Bailion, M.-L. A., A. H. M. van Vliet, J. M. Ketley, C. Constantinidou, and C. W. Penn. 1999. An iron-regulated alkyl hydroperoxide reductase (AhpC) confers aerotolerance and oxidative stress resistance to the microaerophilic pathogen *Campylobacter jejuni*. *J. Bacteriol.* **181**:4798–4804.
- Barriere, C., R. Bruckner, D. Centeno, and R. Talon. 2002. Characterisation of the *katA* gene encoding a catalase and evidence for at least a second catalase activity in *Staphylococcus xylosum*, bacteria used in food fermentation. *FEMS Microbiol. Lett.* **216**:277–283.
- Bender, K. S., H.-C. Yen, and J. D. Wall. 2006. Analysing the metabolic capabilities of *Desulfovibrio* species through genetic manipulation. *Biotechnol. Genet. Eng. Rev.* **23**:157–174.
- Boyd, J., M. N. Oza, and J. R. Murphy. 1990. Molecular cloning and DNA-sequence analysis of a diphtheria *tox* iron-dependent regulatory element (*dxr*) from *Corynebacterium diphtheriae*. *Proc. Natl. Acad. Sci. USA* **87**:5968–5972.
- Brandis, A., and R. K. Thauer. 1981. Growth of *Desulfovibrio* species on hydrogen and sulfate as sole energy source. *J. Gen. Microbiol.* **12**:249–252.
- Brickman, T. J., and S. K. Armstrong. 1995. *Bordetella pertussis fur* gene restores iron repressibility of siderophore and protein expression to deregulated *Bordetella bronchiseptica* mutants. *J. Bacteriol.* **177**:268–270.
- Bsat, N., A. Herbig, L. Casillas-Martínez, P. Setlow, and J. D. Helmann. 1998. *Bacillus subtilis* contains multiple Fur homologues; identification of the iron uptake (Fur) and peroxide regulon (PerR) repressors. *Mol. Microbiol.* **29**:189–198.
- Chao, T.-C., J. Buhrmester, N. Hansmeier, A. Puhler, and S. Weidner. 2005. Role of the regulatory gene *irrA* in the transcriptional response of *Sinorhizobium meliloti* to iron limitation. *Appl. Environ. Microbiol.* **71**:5969–5982.
- Chen, L., L. Keramati, and J. D. Helmann. 1995. Coordinate regulation of *Bacillus subtilis* peroxide stress genes by hydrogen peroxide and metal ions. *Proc. Natl. Acad. Sci. USA* **92**:8190–8194.
- Chhabra, S. R., Q. He, K. H. Huang, S. P. Gaucher, E. J. Alm, Z. He, M. Z. Hadi, T. C. Hazen, J. D. Wall, A. P. Arkin, and A. K. Singh. 2006. Global analysis of heat shock response in *Desulfovibrio vulgaris* Hildenborough. *J. Bacteriol.* **188**:1817–1828.
- Colantuoni, C., G. Henry, S. Zeger, and J. Pevsner. 2002. Local mean normalization of microarray element signal intensities across an array surface: quality control and correction of spatially systematic artifacts. *BioTechniques* **32**:1316–1320.
- Danielli, A., D. Roncarati, I. Delany, V. Chiarini, R. Rappuoli, and V. Scarlato. 2006. In vivo dissection of the *Helicobacter pylori* Fur regulatory circuit by genome-wide location analysis. *J. Bacteriol.* **188**:4654–4662.
- D'Autreaux, B., D. Touati, B. Bersch, J. M. Latour, and I. Michaud-Soret. 2002. Direct inhibition by nitric oxide of the transcriptional ferric uptake regulation protein via nitrosylation of the iron. *Proc. Natl. Acad. Sci. USA* **99**:16619–16624.
- Delany, I., R. Ieva, A. Soragni, M. Hilleringmann, R. Rappuoli, and V. Scarlato. 2005. In vitro analysis of protein-operator interactions of the NikR and Fur metal-responsive regulators of coregulated genes in *Helicobacter pylori*. *J. Bacteriol.* **187**:7703–7715.
- Delany, I., R. Rappuoli, and V. Scarlato. 2004. Fur functions as an activator and as a repressor of putative virulence genes in *Neisseria meningitidis*. *Mol. Microbiol.* **52**:1081–1090.
- Delany, I., G. Spohn, R. Rappuoli, and V. Scarlato. 2001. The Fur repressor controls transcription of iron-activated and -repressed genes in *Helicobacter pylori*. *Mol. Microbiol.* **42**:1297–1309.
- Dos Santos, W. G., I. Pacheco, M. Y. Liu, M. Teixeira, A. V. Xavier, and J. Le Gall. 2000. Purification and characterization of an iron superoxide dismutase and catalase from the sulfate-reducing bacterium *Desulfovibrio gigas*. *J. Bacteriol.* **182**:796–804.
- Elias, D. A., L. R. Krumholz, D. Wong, P. E. Long, and J. M. Sufita. 2003. Characterization of microbial activities and U reduction in a shallow aquifer contaminated by uranium mill tailings. *Microb. Ecol.* **46**:83–91.
- Ernst, F. D., S. Bereswill, B. Waidner, J. Stoeff, U. Mäder, J. G. Kusters, E. J. Kuipers, M. Kist, A. H. M. van Vliet, and G. Homuth. 2005. Transcriptional profiling of *Helicobacter pylori* Fur- and iron-regulated gene expression. *Microbiology* **151**:533–546.
- Ernst, F. D., G. Homuth, J. Stoeff, U. Mäder, B. Waidner, E. J. Kuipers, M. Kist, J. Kusters, S. Bereswill, and A. H. M. van Vliet. 2005. Iron-responsive regulation of the *Helicobacter pylori* iron-cofactored superoxide dismutase SodB is mediated by Fur. *J. Bacteriol.* **187**:3687–3692.
- Escobar, L., J. Perez-Martin, and V. de Lorenzo. 1999. Opening the iron box: transcriptional metalloregulation by the Fur protein. *J. Bacteriol.* **181**:6223–6229.
- Frazaõ, C., G. Silva, C. M. Gomes, P. Matias, R. Coelho, L. Sieker, S. Macedo, M. Y. Liu, S. Oliveira, M. Teixeira, A. V. Xavier, C. Rodrigues-Pousada, M. A. Carrondo, and J. L. Gall. 2000. Structure of a dioxygen reductase enzyme from *Desulfovibrio gigas*. *Nat. Struct. Biol.* **7**:1041–1045.
- Frederick, J. R., J. G. Elkins, N. Bollinger, D. J. Hassett, and T. R. McDermott. 2001. Factors affecting catalase expression in *Pseudomonas aeruginosa* biofilms and planktonic cells. *Appl. Environ. Microbiol.* **67**:1375–1379.
- Fu, R., and G. Voordouw. 1997. Targeted gene-replacement mutagenesis of *dcrA*, encoding an oxygen sensor of the sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Microbiology* **143**:1815–1826.
- Giaever, G., A. M. Chu, L. Ni, C. Connelly, L. Riles, S. Veronneau, S. Dow, A. Lucau-Danila, K. Anderson, B. Andre, A. P. Arkin, A. Astromoff, M. El-Bakkoury, R. Bangham, R. Benito, S. Brachat, S. Campanaro, M. Curtiss, K. Davis, A. Deutschbauer, K. D. Entian, P. Flaherty, F. Foury, D. J. Garfinkel, M. Gerstein, D. Gotte, U. Guldeher, J. H. Hegemann, S. Hempel, Z. Herman, D. F. Jaramillo, D. E. Kelly, S. L. Kelly, P. Kötter, D. LaBonte, D. C. Lamb, N. Lan, H. Liang, H. Liao, L. Liu, C. Luo, M. Lussier, R. Mao, P. Menard, S. L. Ooi, J. L. Revuelta, C. J. Roberts, M. Rose, P. Ross-Macdonald, B. Scherens, G. Schimmack, B. Shafer, D. D. Shoemaker, S. Sookhai-Mahadeo, R. K. Storms, J. N. Strathern, G. Valle, M. Voet, G. Volckaert, C. Y. Wang, T. R. Ward, J. Wilhelmy, E. A. Winzler, Y. Yang, G. Yen, E. Youngman, K. Yu, H. Bussey, J. D. Boeke, M. Snyder, P. Philippsen, R. W. Davis, and M. Johnston. 2002. Functional profiling of the *Saccharomyces cerevisiae* genome. *Nature* **418**:387–391.
- Hamza, I., Z. Qi, N. D. King, and M. R. O'Brian. 2000. Fur-independent regulation of iron metabolism by Irr in *Bradyrhizobium japonicum*. *Microbiology* **146**:669–676.
- Hantke, K. 2001. Iron and metal regulation in bacteria. *Curr. Opin. Microbiol.* **4**:172–177.
- Hantke, K. 1987. Selection procedure for deregulated iron transport mutants (*fur*) in *Escherichia coli* K12: *fur* not only affects iron metabolism. *Mol. Gen. Genet.* **210**:135–139.
- Hantke, K., and V. Braun. 2000. The art of keeping low and high iron concentrations in balance, p. 275–288. In G. Storz and R. Hengge-Aronis (ed.), *Bacterial stress responses*. ASM Press, Washington, D.C.
- Harasthy, V. I., S. F. Jordan, and J. J. Zamboni. 2006. Identification of Fur-regulated genes in *Actinobacillus actinomycetemcomitans*. *Microbiology* **152**:787–796.
- Harris, A. G., F. E. Hinds, A. G. Beckhouse, T. Kolesnikow, and S. L. Hazell. 2002. Resistance to hydrogen peroxide in *Helicobacter pylori*: role of catalase (KatA) and Fur, and functional analysis of a novel gene product designated 'KatA-associated protein', KapA (HP0874). *Microbiology* **148**:3813–3825.
- Hassett, D. J., P. A. Sokol, M. L. Howell, J. F. Ma, H. T. Schweizer, U. Ochsner, and M. L. Vasil. 1996. Ferric uptake regulator (Fur) mutants of *Pseudomonas aeruginosa* demonstrate defective siderophore-mediated iron uptake, altered aerobic growth, and decreased superoxide dismutase and catalase activities. *J. Bacteriol.* **178**:3996–4003.
- He, Q., K. H. Huang, Z. He, E. J. Alm, M. W. Fields, T. C. Hazen, A. P. Arkin, J. D. Wall, and J. Zhou. 2006. Energetic consequences of nitrite stress in *Desulfovibrio vulgaris* Hildenborough, inferred from global transcriptional analysis. *Appl. Environ. Microbiol.* **72**:4370–4381.
- Heidelberg, J. F., R. Seshadri, S. A. Haveman, C. L. Hemme, I. T. Paulsen, J. F. Kolonay, J. A. Eisen, N. Ward, B. Methe, L. M. Brinkac, S. C. Daugherty, R. T. Deboy, R. J. Dodson, A. S. Durkin, R. Madupu, W. C. Nelson, S. A. Sullivan, D. Fouts, D. H. Haft, J. Selengut, J. D. Peterson, T. M. Davidsen, N. Zafar, L. Zhou, D. Radune, G. Dimitrov, M. Hance, K. Tran, H. Khouri, J. Gill, T. R. Utterback, T. V. Feldblyum, J. D. Wall, G. Voordouw, and C. M. Fraser. 2004. The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Nat. Biotechnol.* **22**:1–6.
- Hoffmann, T., A. Schütz, M. Brosius, A. Völker, U. Völker, and E. Bremer. 2002. High-salinity-induced iron limitation in *Bacillus subtilis*. *J. Bacteriol.* **184**:718–727.
- Holmes, K., F. Mulholland, B. M. Pearson, C. Pin, J. McNicholl-Kennedy, J. M. Ketley, and J. M. Wells. 2005. *Campylobacter jejuni* gene expression in response to iron limitation and the role of Fur. *Microbiology* **151**:243–257.
- Köster, W. 2001. ABC transporter-mediated uptake of iron, siderophores, heme, and vitamin B₁₂. *Res. Microbiol.* **152**:291–301.
- Lam, M. S., C. M. Litwin, P. A. Carroll, and B. Calderwood. 1994. *Vibrio cholerae fur* mutations associated with loss of repressor activity: implications for the structural-functional relationships of *fur*. *J. Bacteriol.* **176**:5108–5115.
- Lee, J. W., and J. D. Helmann. 2006. The PerR transcription factor senses H₂O₂ by metal-catalysed histidine oxidation. *Nature* **440**:363–367.
- Londry, K. L., and J. M. Sufita. 1999. Use of nitrate to control sulfide

- generation by sulfate-reducing bacteria associated with oily waste. *J. Ind. Microbiol. Biotechnol.* **22**:582–589.
43. Louvel, H., I. Saint Girons, and M. Picardeau. 2005. Isolation and characterization of FecA- and FecB-mediated iron acquisition systems of the spirochete *Leptospira biflexa* by random insertional mutagenesis. *J. Bacteriol.* **187**:3249–3254.
 44. Moeck, G. S., and J. W. Coulton. 1998. TonB-dependent iron acquisition: mechanisms of siderophore-mediated active transport. *Mol. Microbiol.* **28**: 675–681.
 45. Moura, I., P. Tavares, and N. Ravi. 1994. Characterization of 3 proteins containing multiple iron sites—rubrerythrin, desulfoferrodoxin, and a protein containing a six-iron cluster. *Methods Enzymol.* **243**:216–240.
 46. Mukhopadhyay, A., Z. He, E. J. Alm, A. P. Arkin, E. E. Baidoo, S. C. Borglin, W. Chen, T. C. Hazen, Q. He, H. Y. Holman, K. Huang, R. Huang, D. C. Joyner, N. Katz, M. Keller, P. Oeller, A. Redding, J. Sun, J. D. Wall, J. Wei, Z. Yang, H.-C. Yen, J. Zhou, and J. D. Keasling. 2006. Salt stress in *Desulfovibrio vulgaris* Hildenborough: an integrated genomics approach. *J. Bacteriol.* **188**:4068–4078.
 47. Mukhopadhyay, A., A. M. Redding, M. P. Joachimiak, A. P. Arkin, S. E. Borglin, P. S. Dehal, R. Chakraborty, J. T. Geller, T. C. Hazen, Q. He, D. C. Joyner, V. J. J. Martin, J. D. Wall, Z. K. Yang, J. Zhou, and J. D. Keasling. 2007. Cell wide responses to low oxygen exposure in *Desulfovibrio vulgaris* Hildenborough. *J. Bacteriol.* **189**:5996–6010.
 48. Mukhopadhyay, P., M. Zheng, L. A. Bedzyk, R. A. LaRossa, and G. Storz. 2004. Prominent roles of the NorR and Fur regulators in the *Escherichia coli* transcriptional response to reactive nitrogen species. *Proc. Natl. Acad. Sci. USA* **101**:745–750.
 49. Myhr, S., B. L. P. Lillebo, E. Sunde, J. Beeder, and T. Torsvik. 2002. Inhibition of microbial H₂S production in an oil reservoir model column by nitrate injection. *Appl. Microbiol. Biotechnol.* **58**:400–408.
 50. Nemati, M., T. J. Mazutinec, G. E. Jenneman, and G. Voordouw. 2001. Control of biogenic H₂S production with nitrite and molybdate. *J. Ind. Microbiol. Biotechnol.* **26**:350–355.
 51. Neugebauer, H., C. Herrmann, W. Kammer, G. Schwarz, A. Nordheim, and V. Braun. 2005. ExbBD-dependent transport of maltodextrins through the novel MalA protein across the outer membrane of *Caulobacter crescentus*. *J. Bacteriol.* **187**:8300–8311.
 52. Parker, D., R. M. Kennan, G. S. Myers, I. T. Paulsen, and J. I. Rood. 2005. Identification of a *Dichelobacter nodosus* ferric uptake regulator and determination of its regulatory targets. *J. Bacteriol.* **187**:366–375.
 53. Postgate, J. R. 1984. The sulfate-reducing bacteria, 2nd ed. Cambridge University Press, Cambridge, United Kingdom.
 54. Postle, K., and J. Kadner. 2003. Touch and go: tying TonB to transport. *Mol. Microbiol.* **49**:869–882.
 55. Prince, R. W., C. D. Cox, and M. L. Vasil. 1993. Coordinate regulation of siderophore and exotoxin A production: molecular cloning and sequencing of the *Pseudomonas aeruginosa* fur gene. *J. Bacteriol.* **175**:2589–2598.
 56. Quatrini, R., E. Jedlicki, and D. S. Holmes. 2005. Genomic insights into the iron uptake mechanisms of the biomining microorganism *Acidithiobacillus ferrooxidans*. *J. Ind. Microbiol. Biotechnol.* **32**:606–614.
 57. Rapp-Giles, B. J., L. Casalot, R. S. English, J. A. Ringbauer, Jr., A. Dolla, and J. D. Wall. 2000. Cytochrome c₃ mutants of *Desulfovibrio desulfuricans*. *Appl. Environ. Microbiol.* **66**:671–677.
 58. Ratnayake, D. B., S. N. Wai, Y. Shi, K. Amako, H. Nakayama, and K. Nakayama. 2000. Ferritin from the obligate anaerobe *Porphyromonas gingivalis*: purification, gene cloning and mutant studies. *Microbiology* **146**:1119–1127.
 59. Redding, A. M., A. Mukhopadhyay, D. C. Joyner, T. C. Hazen, and J. D. Keasling. 2006. Study of nitrate stress in *Desulfovibrio vulgaris* Hildenborough using iTRAQ proteomics. *Brief. Funct. Genomics Proteomics* **5**:133–143.
 60. Rocha, E. R., S. C. Andrews, J. N. Keen, and J. H. Brock. 1992. Isolation of a ferritin from *Bacteroides fragilis*. *FEMS Microbiol. Lett.* **74**:207–212.
 61. Rodionov, D. A., I. Dubchak, A. Arkin, E. Alm, and M. S. Gelfand. 2004. Reconstruction of regulatory and metabolic pathways in metal-reducing δ-proteobacteria. *Genome Biol.* **5**:R90.
 62. Romão, C. V., M. Regalla, A. V. Xavier, M. Teixeira, M. Y. Liu, and J. Le Gall. 2000. A bacterioferritin from the strict anaerobe *Desulfovibrio desulfuricans* ATCC 27774. *Biochemistry* **39**:6841–6849.
 63. Rousset, M., L. Casalot, B. J. Rapp-Giles, Z. Dermoun, P. de Philip, J. P. Bélaich, and J. D. Wall. 1998. New shuttle vectors for the introduction of cloned DNA in *Desulfovibrio*. *Plasmid* **39**:114–122.
 64. Rousset, M., Z. Dermoun, M. Chippaux, and J. P. Bélaich. 1991. Marker exchange mutagenesis of the *hydN* genes in *Desulfovibrio fructosovorans*. *Mol. Microbiol.* **5**:1735–1740.
 65. Rowland, B. M., and H. W. Taber. 1996. Duplicate isochorismate synthase genes of *Bacillus subtilis*: regulation and involvement in the biosyntheses of menaquinone and 2,3-dihydroxybenzoate. *J. Bacteriol.* **178**:854–861.
 66. Ryan, R. P., Y. Fouhy, J. F. Lucey, and J. M. Dow. 2006. Cyclic di-GMP signaling in bacteria: recent advances and new puzzles. *J. Bacteriol.* **188**: 8327–8334.
 67. Schalk, I. J., W. W. Yue, and S. K. Buchanan. 2004. Recognition of iron-free siderophores by TonB-dependent iron transporters. *Mol. Microbiol.* **54**: 14–22.
 68. Schneider, R., and K. Hantke. 1993. Iron-hydroxamate uptake systems in *Bacillus subtilis*: identification of a lipoprotein as part of a binding protein-dependent transport system. *Mol. Microbiol.* **8**:111–121.
 69. Shoemaker, D., D. A. Lashkari, D. Morris, M. Mittmann, and R. W. Davis. 1996. Quantitative phenotypic analysis of yeast deletion mutants using a highly parallel molecular bar-coding strategy. *Nat. Genet.* **14**:450–456.
 70. Steil, L., T. Hoffmann, I. Budde, U. Völker, and E. Bremer. 2003. Genome-wide transcriptional profiling analysis of adaptation of *Bacillus subtilis* to high salinity. *J. Bacteriol.* **185**:6358–6370.
 71. Thompson, D. K., A. S. Beliaev, C. S. Giometti, S. L. Tollaksen, T. Khare, D. P. Lies, K. H. Neilson, H. Lim, J. Yates III, C. C. Brandt, and J. M. Tiedje. 2002. Transcriptional and proteomic analysis of a ferric uptake regulator (*fur*) mutant of *Shewanella oneidensis*: possible involvement of *fur* in energy metabolism, transcriptional regulation, and oxidative stress. *Appl. Environ. Microbiol.* **68**:881–892.
 72. Touati, D. 2000. Iron and oxidative stress in bacteria. *Arch. Biochem. Biophys.* **373**:1–6.
 73. van Vliet, A. H., M. L. Baillon, C. W. Penn, and J. M. Ketley. 1999. *Campylobacter jejuni* contains two Fur homologs: characterization of iron-responsive regulation of peroxide stress defense genes by the PerR repressor. *J. Bacteriol.* **181**:6371–6376.
 74. Wall, J. D., H.-C. Yen, and E. C. Drury. 2007. Evaluation of stress response in sulphate-reducing bacteria through genome analysis, p. 141–165. In L. L. Barton (ed.), *Sulphate-reducing bacteria*. Cambridge University Press, Cambridge, United Kingdom.
 75. Wildschut, J. D., R. M. Lang, J. K. Voordouw, and G. Voordouw. 2006. Rubredoxin: oxygen oxidoreductase enhances survival of *Desulfovibrio vulgaris* Hildenborough under microaerophilic conditions. *J. Bacteriol.* **188**: 6253–6260.