

# Real-time molecular monitoring of chemical environment in obligate anaerobes during oxygen adaptive response

Hoi-Ying N. Holman<sup>a,b,1</sup>, Eleanor Wozei<sup>a,2</sup>, Zhang Lin<sup>a,3</sup>, Luis R. Comolli<sup>a</sup>, David A. Ball<sup>a</sup>, Sharon Borglin<sup>a,b</sup>, Matthew W. Fields<sup>b,c</sup>, Terry C. Hazen<sup>a,b</sup>, and Kenneth H. Downing<sup>a</sup>

<sup>a</sup>Lawrence Berkeley National Laboratory, One Cyclotron Road, Berkeley, CA 94720; <sup>b</sup>Virtual Institute for Microbial Stress and Survival, Lawrence Berkeley National Laboratory, Berkeley, CA 94720; and <sup>c</sup>Montana State University, Bozeman, MT 59717

Edited by Jerrold Meinwald, Cornell University, Ithaca, NY, and approved May 12, 2009 (received for review February 25, 2009)

Determining the transient chemical properties of the intracellular environment can elucidate the paths through which a biological system adapts to changes in its environment, for example, the mechanisms that enable some obligate anaerobic bacteria to survive a sudden exposure to oxygen. Here we used high-resolution Fourier transform infrared (FTIR) spectromicroscopy to continuously follow cellular chemistry within living obligate anaerobes by monitoring hydrogen bond structures in their cellular water. We observed a sequence of well orchestrated molecular events that correspond to changes in cellular processes in those cells that survive, but only accumulation of radicals in those that do not. We thereby can interpret the adaptive response in terms of transient intracellular chemistry and link it to oxygen stress and survival. This ability to monitor chemical changes at the molecular level can yield important insights into a wide range of adaptive responses.

*Desulfovibrio* | hydrogen bond | synchrotron FTIR spectromicroscopy | oxygen stress | cellular water

Significant progress has been made at the biochemical and genetic levels in our understanding of how some environmentally and medically important obligate anaerobes can survive temporarily a sudden exposure to oxygen molecules (1–6). However, our understanding at a cellular molecular level of the actual capacity and mechanisms of how these anaerobes survive remains incomplete (7). The cellular chemical environment fundamentally comprises the nexus between external stimuli and internal biochemical regulatory mechanisms—and affects many properties of cellular adaptive response as well. Determining this transient chemical environment *in vivo* is critical for achieving a more coherent understanding of how some obligate anaerobes adapt to the extreme fluctuations in oxygenation. Such knowledge is seldom complete because it is difficult to make *in vivo* molecular measurements without disturbing cells. In almost all previous studies, cellular chemistry of oxygen-stress adaptive response has been determined by measuring intermediate reaction products in cell extracts taken at selected times (8–10); notably, ongoing changes within a live cell are seldom measured directly. The instability of many of the intermediates greatly complicates measurements of cell extracts and their analyses.

Here, we present results of our using the non-invasive synchrotron radiation-based Fourier transform infrared (FTIR) spectromicroscopy approach (11) to determine the cellular chemical environment by continuously monitoring the dynamics of hydrogen bonding in cellular water *in vivo*. More than 70% of the cellular constituents are highly polar water molecules, and their hydrogen bonding is a useful reflection of the cellular chemical environment because it responds “instantaneously” to ions and other species in their surroundings (12, 13). The infrared spectrum of OH stretch vibrations has been widely used to characterize the dynamics of hydrogen-bonding structures in pure water (13–23). These infrared spectroscopy studies have revealed distinct shifts in vibration frequencies and changes in

spectral shapes and intensities induced by the presence of ions and small molecules (e.g., radicals, small organic acids, and hydrogen gas) in water; similar small molecules are expected to be in cellular water during functional metabolism of the oxygen-stress adaptive response.

In this study, we investigated the dynamics of cellular chemical environment in a model oxygen-stress adaptive response system, namely that of the strictly anaerobic sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough during transient exposure to air. Sulfate-reducing bacteria are of particular interest because of their importance in cycling and transformation of essential nutrients and minerals (24, 25) and of their links to different pathogenesis (26, 27) in environments where extreme fluctuations in oxygen concentrations occur. Among sulfate-reducing bacteria, genome sequencing has shown that *D. vulgaris* has developed well-defined protective enzymatic oxygen-defense systems (5, 24). The bacteria can even survive very high levels of oxygen in their natural environment (28, 29), but the mechanism remains elusive. We demonstrate here that molecular information provided by real-time, *in vivo* FTIR measurements of the transient cellular chemical environment is critical for advancing a fundamental understanding of how these obligate anaerobes adapt to extreme changes during air exposure, by providing direct observations of molecular events measured in the same cells over time.

## Results and Discussion

**Identification of *D. vulgaris* Cells That Can Survive Temporarily in Atmospheric Oxygen.** We first conducted microscopic and spectroscopic analyses to establish at a whole cell level the molecular identity of *D. vulgaris* cells that can survive transient exposure to atmospheric oxygen. This identity enabled us to select the appropriate cells for the real-time FTIR measurements of the oxygen-stress survival response. Fluorescence microscopy images of 2 nucleic acid stains show that most stationary-phase (but not exponential-phase) *D. vulgaris* can survive short-term oxygen exposure. Subsequent electron microscopy images reveal that stationary-phase (but not exponential-phase) *D. vulgaris* accumulates polyglucose (Fig. 1*A*) and stores elemental sulfur particles (Fig. 1*B*). Their FTIR spectrum shows distinctly the

Author contributions: H.-Y.N.H., T.C.H., and K.H.D. designed research; H.-Y.N.H., E.W., and Z.L. performed research; H.-Y.N.H., E.W., Z.L., L.R.C., D.A.B., S.B., M.W.F., and K.H.D. contributed new reagents/analytic tools; H.-Y.N.H., E.W., Z.L., L.R.C., D.A.B., S.B., M.W.F., and K.H.D. analyzed data; and H.-Y.N.H., E.W., Z.L., L.R.C., D.A.B., S.B., M.W.F., and T.C.H., K.H.D. wrote the paper.

The authors declare no conflict of interest.


This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

<sup>1</sup>To whom correspondence should be addressed. E-mail: hholman@lbl.gov.

<sup>2</sup>Present address: The Public Health and Environmental Engineering Laboratory, Department of Civil Engineering, Makerere University, P.O. Box 7062, Kampala, Uganda.


<sup>3</sup>Preset address: Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fujian 350002, People's Republic of China.



**Fig. 1.** Microscopic and spectroscopic analyses of *Desulfovibrio vulgaris*. (A) Left, typical infrared absorption spectra of stationary-phase (red) and exponential-phase (blue) *D. vulgaris*; Right, transmission electron microscopy (TEM) images of thin sections poststained by the periodic acid thiosemicarbazide-osmium (PATO) method (51) show intracellular polyglucose granules (arrows) in stationary-phase but not exponential-phase *D. vulgaris*. (B) Left, cryo-electron microscopy (Cryo-EM) image of a stationary-phase cell containing a large, dense ball; Right, energy dispersive X-ray analysis of freeze-dried cells with (blue) and without (red) such structures. Spectra from areas such as marked by the blue circle in the left image show that the particle contains mainly sulfur. (C) Re-growth of *D. vulgaris* after exposure to air. Note the approximate 20-h lag-time (compared to controls). Different colors represent different viability experiments. (D and E) Cryo-EM (Left) and TEM/PATO (Right) images of *D. vulgaris* after exposure to air for hours show changes in cell membranes, variation in periplasmic space, mottled appearance of cell contents and decreased number of polyglucose granules compared to the unexposed cell in Figs. 1 A and B. The frequency of cells showing such alterations compared to those with substantially more damage suggests that these cells were still alive. (Scale bars, 0.2  $\mu\text{m}$ .)

non-glycosidic polyglucose vibration ( $\nu\text{C}-\text{OH}$ ) band between 1,055 and 1,045  $\text{cm}^{-1}$ , and the glycosidic linkage vibration ( $\nu\text{C}-\text{O}-\text{C}$ ) at  $\approx 1,175 \text{ cm}^{-1}$  (30). These cells can survive air exposure for hours and resume growth when returned to anaerobic conditions (Fig. 1C), despite significant changes in cellular structures and contents (Figs. 1 D and E).

**FTIR Measurement and Analysis Considerations.** Mid-infrared photons emitted from the synchrotron at the Advanced Light Source in the Lawrence Berkeley National Laboratory (CA, USA) were focused through a 15- $\mu\text{m}$  aperture onto a monolayer of stationary-phase *D. vulgaris* cells maintained inside an oxygen-free humidified microscope stage chamber (Fig. 2A). In Fig. 2B is an FTIR spectrum typical of small groups of stationary-phase *D. vulgaris* cells, showing well-resolved vibration bands from polyglucose (30) and other biological macromolecules (31) superimposed on the broad continuum absorption features of the aqueous liquid. To minimize inter-experimental uncertainties, only cells that exhibited spectral features within one standard deviation of the mean (Fig. 2B) were selected for the oxygen-stress adaptive response experiments and the controls. We used the spectrally integrated absorption intensity of the polyglucose  $\nu\text{C}-\text{OH}$  band between  $\approx 1,055$  and 1,045  $\text{cm}^{-1}$  to monitor polyglucose degradation. The intensity of the combination band  $\delta\text{OH} + \nu_{\text{L}}\text{HOH}$  at  $\approx 2,100 \text{ cm}^{-1}$  tracks changes in water content because biomolecules typically show very little absorbance in the  $\delta\text{OH} + \nu_{\text{L}}\text{HOH}$  region (31); the intensity of this band represents water concentration in biological samples (32). To detect changes in the hydrogen-bonding structures in cellular water as a measure of transient chemical environment in living *D. vulgaris*, we first derived the FTIR time-difference spectra (relative to the initial state,  $t = 0$ ) in the hydride-OH dominated stretch region between 1,900 and 3,800  $\text{cm}^{-1}$  from the measured real-time FTIR spectra. In each time-difference spectrum, we minimize the




**Fig. 2.** FTIR measurement setup. (A) Schematic description of the experimental measurement setup. An all-reflective optics infrared microscope focuses the interferometer-modulated synchrotron infrared microbeam through a 15- $\mu\text{m}$  aperture onto a monolayer (see Methods) of live *D. vulgaris*. The reflected signals are collected and sent to the detector. The optical density of this thin film is typically 0.05 at the band dominated by the combined water-bending vibration and libration at  $\approx 2,100 \text{ cm}^{-1}$ . This is an equivalent to a water-film of  $\approx 1.5\text{-}\mu\text{m}$  thick. This experimental system enables FTIR measurements with a temporal resolution of every minute for up to 4 h; a different experimental arrangement would be needed to investigate changes on a finer temporal scale or a longer duration. (B) Spectral variations in polyglucose-accumulated stationary-phase *D. vulgaris* in anaerobic atmosphere. The spectrum shows the polyglucose C-OH vibration ( $\nu\text{C}-\text{OH}$ ) band between 1,055 and 1,045  $\text{cm}^{-1}$ . Within the hydride-OH dominated stretch region between 1,900 and 3,800  $\text{cm}^{-1}$  are a broad OH stretching ( $\nu\text{OH}$ ) band between 2,900 and 3,700  $\text{cm}^{-1}$ , the combined water OH bending and libration modes ( $\delta\text{OH} + \nu_{\text{L}}\text{HOH}$ ) at  $\approx 2,100 \text{ cm}^{-1}$ . Absorption bands between 1,800 and 900  $\text{cm}^{-1}$  are dominated by vibration motion of biomolecules of *D. vulgaris*. Averaged spectrum (black line)  $\pm 1.0$  standard deviation (gray lines);  $n = 50$ .

water-continuum absorbance; a positive absorption band reflects the formation of intermediates while a negative band the depletion of an initial state. To interpret changes observed in the FTIR time-difference spectra and to link them to the presence of ions and other small molecules, we used results from previous infrared simulation studies and infrared measurements on aqueous liquid and water clusters (15–23). Because water molecules simultaneously can be hydrogen donors and acceptors, whether the water be liquid or small clusters, spectral information from vibrational spectra of water clusters can be applied to understand dynamics in liquid or other condensed phases (33, 34) such as water in the cellular environment.

**Real-Time FTIR Spectromicroscopy of Oxygen-Stress Adaptive Response in Live *D. vulgaris*.** We monitored cells first under anaerobic conditions and then exposed to air. The measurements are shown in Figs. 3 and 4, respectively, with comparisons shown in Figs. 4C and D. The experiments proceeded as follows. We first made real-time FTIR measurements on a monolayer of *D. vulgaris* inside the oxygen-free humidified microscope stage chamber [hereafter *D. vulgaris*(+polyG;-air)] every 5 min for 240 min without interruption. Fig. 3A shows the real-time FTIR spectra. An easy way to analyze and understand the time-difference spectra is to make a 2-dimensional time-frequency contour plot of the difference spectra in the hydride-OH stretch region, as shown in Fig. 3B (negative values are shown in dark blue) with difference spectrum snapshots below. The plot shows, for  $t < \approx 50$  min, positive bands at  $\approx 3,190 \text{ cm}^{-1}$ , 3,645  $\text{cm}^{-1}$ , and a shoulder feature at 3,745  $\text{cm}^{-1}$ . These frequencies are in the  $\nu\text{OH}$  regions of H-bonding structures of water molecules surrounded by hydrogen gas  $\text{H}_2$  (i.e., hydrogen hydrates) (14). This increasing positive behavior suggests a temporarily enhanced hydrogen gas production event, which is consistent with the central metabolism of *D. vulgaris* under anaerobic conditions (24, 35). This spectral information is taken as reference.





**Fig. 4.** FTIR analyses of *D. vulgaris* during oxygen-stress adaptive response. (A) Typical real-time FTIR spectra of polyglucose-accumulated stationary-phase *D. vulgaris* transition from an anaerobic to aerobic environment. Sequential spectra are offset upward for clarity. Since all spectra are derived using air as a reference, the abrupt change in the spectral feature at  $\approx 2,348\text{ cm}^{-1}$  associated with the presence of atmospheric  $\text{CO}_2$ . (B) Corresponding FTIR time-difference spectra in the hydride-OH dominated stretch region. *Top*, a 2-dimensional frequency-time contour plot (the time-difference intensities are normalized to the maximum); *Bottom*, snapshots for selected different time points. The dashed line marks that difference absorbance = 0. Positive values arise from  $\nu\text{OH}$  of water molecules forming H-bonding with acetate ( $\approx 3,440$ ;  $\approx 2,930\text{ cm}^{-1}$ ) [labeled as  $\nu\text{OH}(\text{water}\cdots\text{acetate})$ ], reactive oxygen species ROS ( $\approx 3,090\text{ cm}^{-1}$ ) [labeled as  $\nu\text{OH}(\text{water}\cdots\text{ROS})$ ], sulfate ( $\approx 3,565\text{ cm}^{-1}$ ) [labeled as  $\nu\text{OH}(\text{water}\cdots\text{sulfate})$ ], and carbonic acid ( $\approx 2,450\text{ cm}^{-1}$ ) [labeled as  $\nu\text{OH}(\text{water}\cdots\text{CO}_2)$ ]. The positive absorption feature at  $\approx 2,360\text{ cm}^{-1}$  is from  $\text{CO}_2$  in air. (C and D). Typical time course of infrared intensity (normalized by the maximum value) of water and polyglucose content. (Bars,  $\pm 10\%$  error.) (E) Transient chemistry as seen by the time-course of difference absorbance normalized by the maximum value for each species. (Bars,  $\pm 10\%$  error.)


pearance of ROS (compare the pink hexagons to the red squares in Fig. 4E) and the increase of water content (green inverted triangles in Fig. 4C).

There is increasing spectral complexity after  $t = \approx 70\text{ min}$ , as suggested by the pattern of the contours in Fig. 4B. The complexity includes progressive band broadening ( $\approx 100\text{ cm}^{-1}$ ) and a redshift ( $\approx 250\text{ cm}^{-1}$ ) in the existing  $\nu\text{OH}$  band of the water $\cdots$ ROS system (between  $\approx 3,200$  and  $3,030\text{ cm}^{-1}$ ) as well as the water $\cdots$  $\text{CO}_2$  system (at frequencies  $< \approx 2,650\text{ cm}^{-1}$ ). This may arise partly from an increase in water molecules available to H-bond with either other anions or neutral species in *D. vulgaris* (+polyG; +air), or other more complicated factors that distort the spectral character (44, 45). For longer times ( $t > \approx 150\text{ min}$ ), the intensity of the  $\nu\text{OH}$  band of the water $\cdots$ ROS declines (red squares in Fig. 4E), which suggests an improved ROS removal in *D. vulgaris* (+polyG; +air).

**Confirmation.** To confirm our results, we therefore used the same method and examined how the FTIR spectra in the hydride-OH dominated stretch region would differ in *D. vulgaris* that had not accumulated intracellular polyglucose granules [hereafter *D. vulgaris* (-polyG; +air)]. Unlike the *D. vulgaris* (+polyG; +air), a positive broad  $\nu\text{OH}$  feature appeared immediately in the frequency range associated with the ionic H-bond with anionic ROS (15–18, 22) (Fig. 5 Upper). The feature broadened by

several hundreds of  $\text{cm}^{-1}$  toward lower frequencies, and reflects an initial disordered ensemble of many different OH stretching vibration modes. The ROS formation continued to exceed their removal with air exposure time, as demonstrated by the monotonically increasing intensity and gradual red-shifts of the vibration modes (centered at  $\approx 3,100\text{ cm}^{-1}$  initially to  $\approx 3,050\text{ cm}^{-1}$  at later times). The immediate positive feature detected at  $\approx 3,570\text{ cm}^{-1}$  (the red trace in Fig. 5 Lower) is within the spectral region of water molecules H-bonded to sulfate anions (23). This small positive value is consistent with our earlier observation that *D. vulgaris* in exponential phase accumulates little elemental sulfur. Little growth was detected when these *D. vulgaris* were returned to oxygen-free conditions.

**Conclusions and Implications.** Our interpretation of the  $\nu\text{OH}$  bands in the FTIR time-difference spectra along with the time-course of the polyglucose  $\nu\text{C-OH}$  and the water ( $\delta\text{OH} + \nu_{\text{L}}\text{HOH}$ ) band intensities is summarized in Fig. 6. Considering the complexity of a living bacterial system, the consistency of the spectral features and the agreement with the putative events of oxygen-stress adaptive response is striking. We have thus demonstrated both the efficacy of using the hydrogen bonding in water of living cells to profile intracellular chemical environment and their significant consequences for understanding functional metabolic controls in “obligate” anaerobic bacteria that can survive oxy-



**Fig. 5.** Typical FTIR difference spectra show reactive oxygen species (ROS) build-up in polyglucose-deficient exponential-phase *D. vulgaris*. *Top*, a 2-dimensional frequency-time contour plot (the time-difference intensities are normalized to the maximum); *Bottom*, snapshots for selected different time points. The peak centered at  $\approx 3,100\text{ cm}^{-1}$  and other local maxima centered at  $\approx 2,904\text{ cm}^{-1}$  and  $2,810\text{ cm}^{-1}$  are at frequencies typical of the  $\nu\text{OH}$  of water molecules H-bonding with hydroxyl and hydroperoxyl radicals [labeled as  $\nu\text{OH}(\text{water}\cdots\text{ROS})$ ]. The feature at  $\approx 3,570\text{ cm}^{-1}$  is at a frequency typical of the  $\nu\text{OH}$  of water molecules H-bonded to sulfate anions. Yellow dots mark the red-shift of  $\approx 75\text{ cm}^{-1}$  of  $\nu\text{OH}$  of hydroxyl radical band peak. The dashed line marks that difference absorbance = 0. The positive absorption feature at  $\approx 2,360\text{ cm}^{-1}$  is from  $\text{CO}_2$  in air.

gen-stress transiently at the chemical level, by providing direct observations of molecular events measured in the same cells over time. Together, these high-resolution synchrotron radiation-based FTIR experiments have revealed a remarkable sequence of well-orchestrated mechanisms that some *D. vulgaris* use to temporarily survive oxygen exposure. When extending this approach to other adaptive-response cellular systems, the experimental design and interpretation of the data should be straightforward in cases where transient chemistry is dominated by ions


or other small chemical species. Even in more complex cases, we anticipate that the interpretation of infrared spectroscopic data in terms of the hydrogen-bonding structure of cellular water will open the door to investigations of chemical and molecular structural changes in living bacteria and other cellular systems over the course of their stress-adaptive response.

## Materials and Methods

**Bacterial Strains and Culture.** *Desulfovibrio vulgaris* Hildenborough (ATCC 29579) was obtained from the American Type Culture Collection. All *D. vulgaris* used in this study were cells of second passage grown on a defined lactate sulfate medium (LS4D medium) soft agar plate (1.0% wt/vol). The LS4D medium was based on Postgate's medium C. To prepare stationary-phase *D. vulgaris* population, we grew *D. vulgaris* (at high cell density) anaerobically on soft LS4D agar until the growth of some bacterial colonies reached confluence. Most cells from the confluent colonies exhibited infrared spectral characteristics typical of polyglucose-containing *D. vulgaris* (red trace in Fig. 1A). To prepare exponential-phase *D. vulgaris* population, we grew *D. vulgaris* (at low cell density) anaerobically on LS4D agar until the growth of bacterial colonies first became visible. Cells from these microcolonies mostly did not exhibit infrared spectral characteristics typical of polyglucose-containing *D. vulgaris* (blue trace in Fig. 1A).

**Preparation of *D. vulgaris* Monolayers.** To ensure that each *D. vulgaris* cell in the FTIR experiment was in contact with atmospheric oxygen, we prepared  $\mu\text{m}$ -thick layers of *D. vulgaris*. We transferred *D. vulgaris* cells onto a LS4D-treated gold-coated glass wafer. An additional mist of liquid LS4D was applied to the replica-printed wafer, which was then incubated for an additional 24 h to facilitate migration of mobile cells to form monolayers on the LS4D-treated gold-coated glass wafer under suitable conditions. To assess the morphological quality of the monolayers, the wafer was placed in a custom microscope stage chamber (filled with nitrogen gas), and was observed by oblique illumination microscopy. Before the FTIR experiment, any excess (moving) LS4D medium was removed by wicking.

**High-Resolution FTIR Spectromicroscopy.** We built a high-humidity microscope stage chamber (Fig. 2A) that allows one to maintain a constant  $\mu\text{m}$ -thick layer of live *D. vulgaris*, overcoming the water interference during high-resolution FTIR spectromicroscopy measurements. All measurements were performed with a Nicolet Magna 760 FTIR bench and a Nicolet Nic-Plan IR microscope equipped with a microscope stage chamber at the infrared beamline of the Advanced Light Source (Lawrence Berkeley National Laboratory, CA, USA; <http://infrared.als.lbl.gov/>). Each spectrum represents an average of 64 scans over a wavenumber range of 4,000 to 650  $\text{cm}^{-1}$  at a spectral resolution of 4  $\text{cm}^{-1}$  with an absorption peak position accuracy of 1/100  $\text{cm}^{-1}$ . As the beam current of the synchrotron decreases with time between electron refills (9 h), the beam intensity decreases proportionally. Appropriate baseline removal took this into account. All data processing was performed using Thermo Electron's Omnic 7.2 (<http://www.thermo.com/>) and Origin 6.0 (<http://www.originlab.com/>). Time-difference spectra are calculated [using Thermo Electron's Omnic 7.2 software (<http://www.thermo.com/>)] from experimental spectra after subtraction of culture medium/liquid water absorbance (46). The time-difference spectra calculation was performed using the water factor approach (47) to minimize absorbance of the water continuum. Analysis of time difference spectra is described in the main text.



**Fig. 6.** A summary of the evolving cellular chemical environment and possible survival mechanisms inside the same living *D. vulgaris* during its transient oxygen-stress and adaptive response, as revealed by the real-time high-resolution synchrotron FTIR measurements and analyses. Polyglucose is labeled as PolyG.

**Carbohydrate Analysis.** Cell pellets were collected from air exposure experiments and were re-suspended in 1 mL of 0.7% NaCl (wt/vol). Protein concentrations were determined with the Lowry assay using BSA (Pierce Biochemicals) as the standard (48). Hexose sugars were measured using the colorimetric cysteine-sulfuric acid method (49) with glucose as the standard.

**Live/Dead Fluorescence Microscopy for Distinguishing Live and Dead Bacteria.** The LIVE/DEAD BacLight bacterial viability kit (Molecular Probes) in combination with a fluorescence microscope (Zeiss) was used. Here, cells were exposed to air for  $t = 0, 30, 240,$  and  $480$  min; and the level of internalized red fluorescing propidium relative to green fluorescing SYTO9 allowed for differentiation between *D. vulgaris* with intact cytoplasmic membranes (bright green) and *D. vulgaris* with damaged cytoplasmic membranes (red/yellow) (50).

**Electron Microscopy.** For conventional TEM/PATO samples, *D. vulgaris* were fixed, embedded, thin sectioned, and post-stained by the periodic acid thiosemicarbazide-osmium (PATO) exactly as previously described (51). Preparation and imaging were conducted at the Robert D. Ogg Electron Microscope Lab at the University of California, Berkeley, CA (<http://em-lab.berkeley.edu/EML/index.php>). Cryo-EM samples were made by placing 5- $\mu$ L aliquots of the *D. vulgaris* suspension onto lacey carbon grids (Ted Ted Pella 01881) that were pretreated by glow-discharge, then blotting and plunging into liquid ethane. Images were recorded on a JEOL-3100 electron microscope operated at 300 kV by a Gatan 795 CCD camera at a magnification of 30,000 $\times$ . Samples to be used

for EDX analysis were freeze-dried in the microscope's airlock and then examined in an FEI CM-200 microscope equipped with a Link EDX detector.

**Re-Growth Experiment.** The re-growth experiments were to obtain growth curves of control (anaerobic) versus air-exposed cells when returned to oxygen-free conditions. Stressed and control cells grown from the oxygen stress experiment were removed from the culture bottles after 24 h of either N<sub>2</sub> or air purge. Ten microliters culture was added to 100  $\mu$ L L54D medium in wells of an empty microplate. Eight replicate wells were used for each culture. Cells were incubated in the Omnilog (Biolog) anaerobically at 30 °C for 100 h. Increase in cell density was quantified by recording increased opacity caused by the accumulation of FeS precipitates. The Omnilog uses a scanning technology, which is analogous to a turbidity measurement, to record the increase in density in Omnilog (OL) units.

**ACKNOWLEDGMENTS.** We thank Dr. K. McDonald and Ms. R. Zalpuri at the Robert D. Ogg Electron Microscope Lab, UC Berkeley, and Dr. Z. Lee at the National Center for Electron Microscopy, Lawrence Berkeley National Laboratory, for technical assistance; Ms. D. Joyner for technical assistance; Drs. H. Bechtel, Z. Hao, J. Jansson, C. Jansson, M.C. Martin, W. McKinney and J. Zhou for discussion and comments on this work and the manuscript. This work was supported by the U.S. Department of Energy Office of Biological and Environmental Research's Structural Biology Program, and Genomics:GTL Program through contract DE-AC02-05CH11231 with Lawrence Berkeley National Laboratory.

- Fay P (1992) Oxygen relations of nitrogen-fixation in Cyanobacteria. *Microbiol Rev* 56:340-373.
- Bradshaw DJ, Marsh PD, Watson GK, Allison C (1998) Role of *Fusobacterium nucleatum* and coaggregation in anaerobe survival in planktonic and biofilm oral microbial communities during aeration. *Infect Immun* 66:4729-4732.
- Cypionka H (2000) Oxygen respiration by *Desulfovibrio* species. *Annu Rev Microbiol* 54:827-848.
- Baughn AD, Malamy MH (2004) Molecular Basis for Aerotolerance of the Obligately Anaerobic *Bacteroides* Spp. In *Strict and Facultative Anaerobes: Medical and Environmental Aspects*, eds Nakano MM, Zuber P (CRC Press, New York), pp 161-169.
- Dolla A, Fournier M, Dermoun Z (2006) Oxygen defense in sulfate-reducing bacteria. *J Biotechnol* 126:87-100.
- Brioukhanov AL, Netrusov AI (2007) Aerotolerance of strictly anaerobic microorganisms and factors of defense against oxidative stress: A review. *Appl Biochem Micro* 43:567-582.
- Muyzer G, Stams AJM (2008) The ecology and biotechnology of sulphate-reducing bacteria. *Nat Rev Microbiol* 6:441-454.
- Fuseler K, Cypionka H (1995) Elemental sulfur as an intermediate of sulfide oxidation with oxygen by *Desulfovibrio propionicus*. *Arch Microbiol* 164:104-109.
- Van Niel EWJ, et al. (1996) The role of polyglucose in oxygen-dependent respiration by a new strain of *Desulfovibrio salexigens*. *Fems Microbiol Ecol* 21:243-253.
- Fareleira P, et al. (2003) Response of a strict anaerobe to oxygen: Survival strategies in *Desulfovibrio gigas*. *Microbiology* 149:1513-1522.
- Holman H-YN, Martin MC (2006) Synchrotron radiation infrared spectrometry: a non-invasive molecular probe for biogeochemical processes. In *Advances in Agronomy*, ed Sparks D (Elsevier, New York), Vol 90, pp 79-127.
- Woutersen S, Bakker HJ (1999) Resonant intermolecular transfer of vibrational energy in liquid water. *Nature* 402:507-509.
- Cowan ML, et al. (2005) Ultrafast memory loss and energy redistribution in the hydrogen bond network of liquid H<sub>2</sub>O. *Nature* 434:199-202.
- Okumura M, Yeh LJ, Myers JD, Lee YT (1990) Infrared-spectra of the solvated hydrogen ion - vibrational predissociation spectroscopy of mass-selected H<sub>3</sub>O<sup>+</sup>. (H<sub>2</sub>O)<sub>n</sub>. (H<sub>2</sub>)<sub>m</sub>. *J Phys Chem-U S* 94:3416-3427.
- Nelander B (1997) The peroxy radical as hydrogen bond donor and hydrogen bond acceptor: A matrix isolation study. *J Phys Chem A* 101:9092-9096.
- Weber JM, et al. (2000) Isolating the spectroscopic signature of a hydration shell with the use of clusters: Superoxide tetrahydrate. *Science* 287:2461-2463.
- Chaudhuri C, et al. (2001) Infrared spectra and isomeric structures of hydroxide ion-water clusters OH<sup>-</sup> (H<sub>2</sub>O)<sub>(n-5)</sub>: A comparison with H<sub>3</sub>O<sup>+</sup> (H<sub>2</sub>O)<sub>(n-5)</sub>. *Mol Phys* 99:1161-1173.
- Robertson WH, et al. (2003) Spectroscopic determination of the OH<sup>-</sup> solvation shell in the OH<sup>-</sup>. (H<sub>2</sub>O)<sub>(n)</sub> clusters. *Science* 299:1367-1372.
- Max JJ, Chapados C (2004) Infrared spectroscopy of aqueous carboxylic acids: Comparison between different acids and their salts. *J Phys Chem A* 108:3324-3337.
- Shin JW, et al. (2004) Infrared signature of structures associated with the H<sup>+</sup> (H<sub>2</sub>O)<sub>(n)</sub> (n=6 to 27) clusters. *Science* 304:1137-1140.
- Headrick JM, et al. (2005) Spectral signatures of hydrated proton vibrations in water clusters. *Science* 308:1765-1769.
- Iyengar SS (2005) Dynamical effects on vibrational and electronic spectra of hydroperoxy radical water clusters. *J Chem Phys* 123:084310-084311.
- Bush MF, Saykally RJ, Williams ER (2007) Evidence for water rings in the hexahydrated sulfide dianion from IR spectroscopy. *J Am Chem Soc* 129:2220-2223.
- Heidelberger JF, et al. (2004) The genome sequence of the anaerobic, sulfate-reducing bacterium *Desulfovibrio vulgaris* Hildenborough. *Nat Biotechnol* 22:554-559.
- Baumgartner LK, et al. (2006) Sulfate reducing bacteria in microbial mats: Changing paradigms, new discoveries. *Sediment Geol* 185:131-145.
- Chapple ILC (1996) Role of free radicals and antioxidants in the pathogenesis of the inflammatory periodontal diseases. *J Clin Pathol-CI Mol* 49:M247-M255.
- Duffy M, et al. (2002) Sulfate-reducing bacteria colonize pouches formed for ulcerative colitis but not for familial adenomatous polyposis. *Dis Colon Rectum* 45:384-388.
- Risatti JB, Capman WC, Stahl DA (1994) Community structure of a microbial mat - the phylogenetic dimension. *P Natl Acad Sci USA* 91:10173-10177.
- Minz D, et al. (1999) Diversity of sulfate-reducing bacteria in oxic and anoxic regions of a microbial mat characterized by comparative analysis of dissimilatory sulfite reductase genes. *Appl Environ Microb* 65:4666-4671.
- Stams FJM, Veenhuis M, Weenk GH, Hansen TA (1983) Occurrence of polyglucose as a storage polymer in *desulfovibrio* species and *desulfovibrio-propionicus*. *Arch Microbiol* 136:54-59.
- Naumann D (2000) Infrared Spectroscopy in Microbiology. In *Encyclopedia of Anal Chem*, ed Meyers RA (John Wiley & Sons Ltd, Chichester), pp 102-131.
- Potts RO, Guzek DB, Harris RR, Mckie JE (1985) A non-invasive, in vivo technique to quantitatively measure water concentration of the stratum-corneum using attenuated total-reflectance infrared-spectroscopy. *Arch Dermatol Res* 277:489-495.
- Vernon MF, et al. (1982) Infrared vibrational predissociation spectroscopy of water clusters by the crossed laser-molecular beam technique. *J Chem Phys* 77:47-57.
- Castleman AW, Keesee RG (1986) Clusters - bridging the gas and condensed phases. *Acc Chem Res* 19:413-419.
- Tsuji K, Yagi T (1980) Significance of hydrogen burst from growing cultures of *Desulfovibrio vulgaris*, Miyazaki, and the role of hydrogenase and cytochrome c<sub>3</sub> in energy-production system. *Arch Microbiol* 125:35-42.
- Baumgarten A, Redenius I, Kranczoch J, Cypionka H (2001) Periplasmic oxygen reduction by *Desulfovibrio* species. *Arch Microbiol* 176:306-309.
- Fitz RM, Cypionka H (1991) Generation of a proton gradient in *Desulfovibrio vulgaris*. *Arch Microbiol* 155:444-448.
- Santos H, et al. (1993) Aerobic metabolism of carbon reserves by the obligate anaerobe *Desulfovibrio gigas*. *Biochem Biophys Res Commun* 195:551-557.
- Brucato JR, Palumbo ME, Strazzulla G (1997) Carbonic acid by ion implantation in water/carbon dioxide ice mixtures. *Icarus* 125:135-144.
- Schauder R, et al. (1986) Acetate oxidation to CO<sub>2</sub> in anaerobic bacteria via a novel pathway not involving reactions of the citric-acid cycle. *Arch Microbiol* 145:162-172.
- Tang Y, et al. (2007) Pathway confirmation and flux analysis of central metabolic pathways in *Desulfovibrio vulgaris* hildenborough using gas chromatography-mass spectrometry and Fourier transform-ion cyclotron resonance mass spectrometry. *J Bacteriol* 189:940-949.
- Kramer M, Cypionka H (1989) Sulfate formation via ATP sulfurylase in thiosulfate-disproportionating and sulfite-disproportionating bacteria. *Arch Microbiol* 151:232-237.
- Fuseler K, Krekeler D, Sydow U, Cypionka H (1996) A common pathway of sulfide oxidation by sulfate-reducing bacteria. *Fems Microbiol Lett* 144:129-134.
- Lawrence CP, Skinner JL (2003) Vibrational spectroscopy of HOD in liquid D<sub>2</sub>O. III. Spectral diffusion, and hydrogen-bonding and rotational dynamics. *J Chem Phys* 118:264-272.
- Yamashita T, Takatsuka K (2007) Hydrogen-bond assisted enormous broadening of infrared spectra of phenol-water cationic cluster: An *ab initio* mixed quantum-classical study. *J Chem Phys* 126:074304.
- Mantele W (1993) Reaction-induced infrared difference spectroscopy for the study of protein function and reaction-mechanisms. *Trends Biochem Sci* 18:197-202.
- Max JJ, Chapados C (2002) Infrared spectroscopy of aqueous carboxylic acid: Malic acid. *J Phys Chem A* 106:6452-6461.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ (1951) Protein measurement with the folin phenol reagent. *J Biol Chem* 193:265-275.
- Chaplin MF (1986) Monosaccharides. In *Carbohydrate analysis*, eds Chaplin, MF, Kennedy JF (IRL Press, Oxford), pp 1-4.
- Leuko S, Legat A, Fendrihan S, Stan-Lotter H (2004) Evaluation of the LIVE/DEAD BacLight kit for detection of extremophilic archaea and visualization of microorganisms in environmental hypersaline samples. *Appl Environ Microb* 70:6884-6886.
- Kropinski AM, Ghiorse WC, Greenberg EP (1988) The intracellular polyglucose storage granules of *Spirochaeta aurantia*. *Arch Microbiol* 150:289-295.